arXiv:hep-th/9810027v1 5 Oct 1998

Two-dimensional Quantum Gravity

Juri Rolf
Niels Bohr Institute
University of Copenhagen

Thesis submitted for the Ph.D. degree in physics
at the Faculty of Science, University of Copenhagen.
May 1998

Contents

Acknowledgement iii

Introduction 1

1 Two-dimensional quantum gravity 2
1.1 Scaling in quantum gravity 3
1.1.1 Partition function 3
1.1.2 Hartle-Hawking wavefunctionals 6
1.1.3 The two-point function 7
1.2 Liouville theory: A brief reminder 11
1.2.1 Functional measures 11
1.2.2 Factorization of the diffeomorphisms 14
1.2.3 Gravitational dressing of scaling exponents 16
1.3 Fractal dimensions 20
1.3.1 Extrinsic Hausdorff dimension 21
1.3.2 Spectral dimension 23
1.3.3 Intrinsic Hausdorff dimension 28

2 Discretization of 2d quantum gravity 32
2.1 Dynamical triangulation 33
2.1.1 Discretization of geometry 33
2.1.2 Definition of the model 36
2.1.3 The two-point function 39
2.1.4 Branched polymers 43
2.2 Matrix models 46
2.2.1 Dynamical triangulation by matrix models 46
2.2.2 The loop equations 49
2.2.3 Scaling limit and computation of γ 52
2.3 The fractal structure of pure gravity 54
2.3.1 The geodesic two-loop function 54

2.3.2 Scaling of the two-point function 55
2.4 Conclusion 57

3 The failure of quantum Regge calculus 58
3.1 Formulation of quantum Regge calculus 59
3.2 Regge integration measures 60
3.2.1 DeWitt-like measures 61
3.2.2 The DeWitt-like measure in other dimensions than two 63
3.2.3 The DeWitt-like measures for special geometries . . . 64
3.2.4 Commonly used measures 65
3.2.5 Numerical simulations in two-dimensional quantum Regge calculus 67
3.3 The appearance of spikes 69
3.3.1 General proof of appearance 70
3.3.2 Spikes in special geometries 72
3.4 Concluding remarks 75
3.4.1 The appearance of spikes in higher dimensions 75
3.4.2 The role of diffeomorphisms in two dimensions 76
3.4.3 Conclusion 79

Conclusion 82

Bibliography 83

Acknowledgement

I want to thank all the people who have directly or indirectly contributed to this work and made my stay in Copenhagen possible, in particular:

- my supervisor Jan Ambjørn for the interesting project, for many discussions, for a good collaboration and for partial financial support of my visit in Santa Fe at the workshop New Directions in Simplicial Quantum Gravity,
- Jakob L. Nielsen for many discussions and for a good collaboration,
- my collaborators Dimitrij Boulatov, George Savvidy and Yoshiyuki Watabiki for the good work done together,
- Martin Harris and Jakob L. Nielsen for proofreading the thesis,
- my parents and my wife for their support and encouragement,
- Konstantinos N. Anagnostopoulos, Martin Harris, Lars Jensen, Jakob L. Nielsen, Kaspar Olsen, Jørgen Rasmussen and Morten Weis for some discussions and for creating the nice atmosphere in our office,
- the Niels Bohr Institute for warm hospitality,
- the Studienstiftung des deutschen Volkes for financial support.

Introduction

This Ph.D. thesis pursues two goals: The study of the geometrical structure of two-dimensional quantum gravity and in particular its fractal nature. To address these questions we review the continuum formalism of quantum gravity with special focus on the scaling properties of the theory. We discuss several concepts of fractal dimensions which characterize the extrinsic and intrinsic geometry of quantum gravity. This work is partly based on work done in collaboration with Jan Ambjørn, Dimitrij Boulatov, Jakob L. Nielsen and Yoshiyuki Watabiki [1].

The other goal is the discussion of the discretization of quantum gravity and to address the so called quantum failure of Regge calculus. We review dynamical triangulations and show that it agrees with the continuum theory in two dimensions. Then we discuss Regge calculus and prove that a continuum limit cannot be taken in a sensible way and that it does not reproduce continuum results. This work is partly based on work done in collaboration with Jan Ambjørn, Jakob L. Nielsen and George Savvidy [2].

In chapter 1 we introduce the main ingredients for the formulation of two-dimensional quantum gravity as an Euclidean functional integral over geometries. It contains a brief reminder of Liouville theory and the technical issues in the continuum formalism. We use these techniques to discuss the extrinsic and intrinsic Hausdorff dimension and the spectral dimension of two-dimensional quantum gravity.

Chapter 2 is a review of dynamical triangulation in two dimensions. We begin with an introduction of the main ideas of how to discretize two-dimensional quantum geometries. The scaling properties are illustrated by means of the two-point function of pure gravity and of branched polymers.

In chapter 3 we discuss quantum Regge calculus which has been suggested as an alternative method to discretize quantum geometries. We prove by a simple scaling argument that a sensible continuum limit of this theory cannot be taken and that it disagrees with continuum results.

Chapter 1

Two-dimensional quantum gravity

Euclidean quantum gravity is an attempt to quantize general relativity based on Feynman's functional integral and on the Einstein-Hilbert action principle. One integrates over all Riemannian metrics on a d-dimensional manifold M. It is based on the hope that one can recover the Lorentzian signature after performing the integration analogously to the Wick rotation in Euclidean quantum field theory. For a general discussion of further problems and for motivation of a theory of quantum gravity we refer to [3, 4, 5].

General relativity is a reparametrization invariant theory which can be formulated with no reference to coordinates at all. This diffeomorphism invariance is a central issue in the quantum theory. Its importance is most apparent in two dimensions, since the Einstein-Hilbert action is trivial and consists only of a topological term and the cosmological constant coupled to the volume of the spacetime. All the non-trivial dynamics of the two-dimensional theory of quantum gravity thus come from gauge fixing the diffeomorphisms while keeping the geometry exactly fixed. This is the famous representation of the functional integral over geometries as a Liouville field theory by Polyakov [6]. Based on this formulation the scaling exponents can be obtained [7, 8, 9].

Any theory of quantum gravity must aim at answering questions about the geometrical structure of quantum spacetime. The interplay between matter and geometry is well known from general relativity. The quantum average over all geometries changes the dynamics of this interaction. It turns out that the quantum spacetime has a fractal nature and even its dimension is a dynamical quantity. The characterization of the fractal nature of the quantum spacetime in two-dimensional quantum gravity is one of the central themes of this work.

In section 1.1 the main concepts in the continuum formalism are intro-

duced. In the following section we review technical details about the factorization of the diffeomorphisms from the functional integral. In section 1.3 we introduce fractal dimensions to characterize the fractal nature of quantum spacetime. That section, and in particular section 1.3.2 are partly based on work presented in [1, 10].

Reviews about the continuum approach to two-dimensional quantum gravity can be found in [11] and the references therein. For an introduction to conformal field theory see [12, 13, 14].

1.1 Scaling in quantum gravity

In this section we introduce the basic concepts in two-dimensional quantum gravity. We begin with the partition function and the Hartle-Hawking wavefunctionals. The most natural object to address questions about the scaling behaviour of the theory is the geodesic two-point function. We discuss its scaling behaviour in detail and introduce the concept of the intrinsic fractal dimension which illustrates the fractal nature of the intrinsic geometry of the two-dimensional quantum space-time.

1.1.1 Partition function

Let M be a two-dimensional, closed, compact, connected, orientable manifold of genus h. Then the partition function for two-dimensional quantum gravity can formally be written as the functional integral

$$Z(G,\Lambda) = \sum_{h\geq 0} \int \mathcal{D}[g_{\mu\nu}]\; e^{-S_{EH}(g;G,\Lambda)} \int \mathcal{D}_g X\; e^{-S_{matter}(g,X)}. \tag{1.1}$$

Here

$$S_{EH}(g;G,\Lambda) = \Lambda \int_M d^2\xi\sqrt{g} - \frac{1}{4\pi G}\int_M d^2\xi\sqrt{g}\;\mathcal{R}(\xi) \tag{1.2}$$

is the classical reparametrization invariant Einstein-Hilbert action [15, 16] with the gravitational coupling constant G, the cosmological constant Λ and the curvature scalar $\mathcal{R}$. According to the Gauss–Bonnet theorem, the last term in (1.2) is a topological invariant, called the Euler characteristic χ of M:

$$\chi(h) = \frac{1}{4\pi}\int_M d^2\xi\sqrt{g}\;\mathcal{R}(\xi) = 2-2h, \tag{1.3}$$

while the first term is the volume V_g of M equipped with the metric $g_{\mu\nu}$:

$$V_g = \int_M d^2\xi \sqrt{g}. \tag{1.4}$$

Therefore, (1.1) can be rewritten as

$$Z(G, \Lambda) = \sum_{h \geq 0} e^{\frac{\chi(h)}{G}} Z(\Lambda), \tag{1.5}$$

where $Z(\Lambda)$ is defined as

$$Z(\Lambda) = \int \mathcal{D}[g_{\mu\nu}]\ e^{-S(g,\Lambda)} \int \mathcal{D}_g X\ e^{-S_{\text{matter}}(g,X)}, \tag{1.6}$$

with $S(g, \Lambda) = \Lambda V_g$.

$S_{\text{matter}}(g, X)$ in (1.1) denotes any reparametrization invariant action for conformal matter fields X with central charge D coupled to gravity. A typical example is the coupling of D free scalar matter fields $X^1, \ldots, X^D$ to gravity. In this case

$$S_{\text{matter}}(g, X) = \frac{1}{8\pi} \int_M d^2\xi \sqrt{g}\ g^{\mu\nu} \partial_\mu X^a \partial_\nu X^a, \tag{1.7}$$

which is diffeomorphism invariant and invariant under Weyl rescalings of the metric:

$$S_{\text{matter}}(e^\phi g, X) = S_{\text{matter}}(g, X). \tag{1.8}$$

Note that (1.7) can also be interpreted as an embedding of M in a D-dimensional Euclidean space, thus leading to an interpretation of (1.1) as bosonic string theory in D dimensions [6].

The functional integration $\int \mathcal{D}[g_{\mu\nu}]$ in (1.1) is an integration over all geometries, that means all diffeomorphism classes $[g_{\mu\nu}]$ of metrics $g_{\mu\nu}$ on the manifold M. This is often denoted formally as

$$\int \frac{\mathcal{D} g_{\mu\nu}}{\text{Vol(Diff)}}, \tag{1.9}$$

where Vol(Diff) is the volume of the group of diffeomorphisms on M. Since this group is not compact the quotient (1.9) does not make sense beyond a formal level. We will derive expressions for the measures $\mathcal{D} g_{\mu\nu}$, $\mathcal{D}[g_{\mu\nu}]$ and $\mathcal{D}_g X$ in section 1.2.

In (1.1) the sum goes over all topologies of two-dimensional manifolds, that means over all genera h. It is presently unknown how to define such a

sum over topologies different from a perturbative expansion in $\hbar$. It comes as a bitter aspect of the theory that we only know how to perform the functional integration in (1.1) for a given manifold and thus only for a given fixed topology, while a summation over topologies has to be performed by hand. In four dimensions it is known that the topologies even cannot be classified. Therefore the role of topological fluctuations is still unclear in two and in higher dimensions. In the remainder of this work we will always fix the topology and thereby disregard the sum over the genera in (1.1).

Sometimes it is useful to define the partition function in an ensemble of universes with fixed volume V:

$$\begin{aligned} Z(V) &= \int \mathcal{D}[g_{\mu\nu}]\ \delta(V - V_g) \int \mathcal{D}_g X\ e^{-S_{\text{matter}}(g,X)} \\ &\equiv \int \mathcal{D}[g_{\mu\nu}]_V \int \mathcal{D}_g X\ e^{-S_{\text{matter}}(g,X)}. \end{aligned} \tag{1.10}$$

$Z(\Lambda)$ is the Laplace transform of $Z(V)$ [1], that means:

$$Z(\Lambda) = \int_0^\infty dV\ e^{-\Lambda V} Z(V), \tag{1.11}$$

$$Z(V) = \int_{\sigma - i\infty}^{\sigma + i\infty} \frac{d\Lambda}{2\pi i}\ e^{\Lambda V} Z(\Lambda), \tag{1.12}$$

where σ is a real constant which exceeds the real part of all the singularities of $Z(\Lambda)$. Formula (1.10) shows that although the action of pure two-dimensional quantum gravity is trivial and does not contain any propagating degrees of freedom, quantum gravity in two dimensions is a true quantum problem, since each equivalence class of metrics is counted once with the same weight. That means that there is no classical spacetime around which we expand and the theory is as "quantum-like" as it can get. Therefore it might contain important information about quantum gravity in higher dimensions.

As we show in section 1.2 $Z(V)$ scales as

$$Z(V) \propto V^{\gamma - 3}, \quad \text{or } Z(\Lambda) \propto \Lambda^{2-\gamma}, \tag{1.13}$$

which defines a scaling exponent γ that depends on the matter coupled to gravity and on the topology of the universe. The computation of γ gives the result [7, 8, 9]

$$\gamma = 2 + \frac{1-h}{12}\left(D - 25 - \sqrt{(25-D)(1-D)}\right). \tag{1.14}$$

[1] That is also the reason why we denote both with the same symbol and distinguish between them by the names of their arguments.

This formula is often called KPZ-formula. For pure gravity ($D = 0$) γ equals $-1/2$, for the Ising model coupled to gravity ($D = 1/2$) we have $\gamma = -1/3$ and at $D = 1$, γ equals 0 for spherical topology. Clearly this formula breaks down for conformal charges $D > 1$, where γ assumes complex values. This has sometimes been called the $D = 1$ barrier. There is now some evidence for the fact that two-dimensional quantum gravity coupled to matter with central charge $D > 1$ is in a branched polymer phase [17, 18] in which the surfaces collapse to tree-like objects.

The expectation value of some observable $\mathcal{O}(g, X)$ is defined as

$$\langle \mathcal{O}(g, X) \rangle_\Lambda = \frac{1}{Z(\Lambda)} \int \mathcal{D}[g_{\mu\nu}] \; e^{-\Lambda V_g} \int \mathcal{D}_g X \; e^{-S_{\text{matter}}(g,X)} \mathcal{O}(g, X), \tag{1.15}$$

or for fixed volume V as

$$\langle \mathcal{O}(g, X) \rangle_V = \frac{1}{Z(V)} \int \mathcal{D}[g_{\mu\nu}]_V \int \mathcal{D}_g X \; e^{-S_{\text{matter}}(g,X)} \mathcal{O}(g, X). \tag{1.16}$$

1.1.2 Hartle-Hawking wavefunctionals

Let us in the remainder of this section ignore the coupling of possible matter fields to quantum gravity. The definitions below can easily be expanded to the general case. Furthermore we concentrate on spherical topology only. Again the generalization is straightforward.

Typical observables in two-dimensional quantum gravity are loop amplitudes, that means amplitudes for one-dimensional universes. Let M be topologically equivalent to a sphere with b holes. The induced gravity on the boundary is covered by the modified action

$$S(g, \Lambda, Z_1, \ldots, Z_b) = \Lambda V_g + \sum_{i=1}^{b} Z_i L_{g,i}, \tag{1.17}$$

where $L_{g,i}$ denotes the length of the i'th boundary component in the metric $g_{\mu\nu}$. In this case the partition function is given by

$$W(\Lambda, Z_1, \ldots, Z_b) = \int \mathcal{D}[g_{\mu\nu}] \; e^{-S(g,\Lambda,Z_1,\ldots,Z_b)}. \tag{1.18}$$

This can also be interpreted as the amplitude for b one-dimensional universes of arbitrary lengths. Since the lengths of the boundary components are invariant under diffeomorphisms it makes sense to fix them to prescribed values $L_1, \ldots, L_b$. This yields the definition of the Hartle-Hawking wave

functionals [19]:

$$W(\Lambda, L_1, \ldots, L_b) = \int \mathcal{D}[g_{\mu\nu}]\ e^{-S(g,\Lambda)} \prod_{i=1}^{b} \delta(L_i - L_{g,i}). \tag{1.19}$$

We note that (1.18) is the Laplace transform of (1.19), that means:

$$W(\Lambda, Z_1, \ldots, Z_b) = \int_0^\infty \prod_{i=1}^{b} dL_i e^{-Z_i L_i}\ W(\Lambda, L_1, \ldots, L_b). \tag{1.20}$$

In the ensemble of universes with fixed volume V the wave functionals are given by:

$$W(V, L_1, \ldots, L_b) = \int \mathcal{D}[g_{\mu\nu}]\ \delta(V - V_g) \prod_{i=1}^{b} \delta(L_i - L_{g,i}). \tag{1.21}$$

The cosmological constant Λ and the volume V are conjugate variables, that means:

$$W(\Lambda, L_1, \ldots, L_b) = \int_0^\infty dV\ e^{-\Lambda V} W(V, L_1, \ldots, L_b). \tag{1.22}$$

In the case $b = 0$ one recovers the partition functions (1.6) and (1.10).

1.1.3 The two-point function

To define reparametrization invariant correlation functions let $d_g(\xi, \xi')$ be the geodesic distance between two points ξ and ξ' with respect to the metric $g_{\mu\nu}$. Then the invariant two-point function is defined as

$$G(\Lambda, R) = \int \mathcal{D}[g_{\mu\nu}]\ e^{-S(g,\Lambda)} \int_M d^2\xi \sqrt{g(\xi)} \int_M d^2\xi' \sqrt{g(\xi')}\ \delta(R - d_g(\xi, \xi')). \tag{1.23}$$

$G(\Lambda, R)$ can also be interpreted as the partition function for universes with two marked points separated by a geodesic distance R. The integrated two-point function, called the susceptibility χ has the following behaviour:

$$\chi(\Lambda) = \int_0^\infty dR\ G(\Lambda, R) = \frac{\partial^2 Z(\Lambda)}{\partial \Lambda^2} \sim \Lambda^{-\gamma}. \tag{1.24}$$

Therefore γ is often called the susceptibility exponent. Yet another characterization of the exponent γ from the branching ratio of the two-dimensional

universes will be explained in chapter 2. For fixed volume of spacetime the definition of the two-point function is

$$G(V,R) = \int \mathcal{D}[g_{\mu\nu}]_V \int_M d^2\xi \sqrt{g(\xi)} \int_M d^2\xi' \sqrt{g(\xi')}\ \delta(R - d_g(\xi,\xi')), \quad (1.25)$$

which is related to formula (1.23) via

$$G(\Lambda,R) = \int_0^\infty dV\ e^{-\Lambda V} G(V,R). \quad (1.26)$$

Similar to (1.23) a whole set of invariant correlation functions which depend on the geodesic distance can be defined by multiplying the local measures $d^2\xi\sqrt{g(\xi)}$ and $d^2\xi'\sqrt{g(\xi')}$ by powers of the invariant curvature scalars $\mathcal{R}(\xi)$ and $\mathcal{R}(\xi')$.

Note that by (1.23) or (1.25) the concept of geodesic distance becomes meaningful even in quantum gravity. The two-point function can be interpreted as the partition function of the ensemble of universes which have two marked points separated by a geodesic distance R. Both, its short distance behaviour and its long distance behaviour reveal the fractal structure of the most important metrics that contribute to the functional integral (1.1).

An important property of the two-point function (1.23) is the inequality

$$G(\Lambda, R_1 + R_2) \geq \text{const} \times G(\Lambda, R_1)\ G(\Lambda, R_2), \quad (1.27)$$

which follows from a simple gluing argument which is explained in chapter 2. Up to a constant, (1.27) is equivalent to the subadditivity of $-\log G(\Lambda, R)$, from which the existence of the limit

$$\lim_{R\to\infty} \frac{-\log G(\Lambda,R)}{R} = M(\Lambda) \quad (1.28)$$

follows from general arguments (see for example [20]). Furthermore one can deduce that $M(\Lambda)$ is an increasing function of Λ and $M(\Lambda) > 0$ for $\Lambda > 0$. At this stage one cannot prove that $M(\Lambda)$ scales to zero as Λ goes to zero. However let us assume that this is the case and verify it later by an explicit calculation of the two-point function in section 2.3:

$$M(\Lambda) = c\ \Lambda^\gamma, \quad (1.29)$$

with a dimensionless constant c. That means that for large $R \gg M(\Lambda)^{-1}$ the two point function falls off exponentially with a subleading correction:

$$G(\Lambda,R) \sim \Lambda^{\nu-\gamma}\ e^{-c\Lambda^\gamma R}, \text{ for } R \gg \frac{1}{M(\Lambda)}. \quad (1.30)$$

The power correction can be found by applying dimensional arguments to (1.24). We conclude that the average volume of two-dimensional universes with two marked points separated by a geodesic distance R is proportional to R if R is large enough:

$$\langle V \rangle_R \equiv -\frac{\partial \log G(\Lambda, R)}{\partial \Lambda} \underset{R \gg M(\Lambda)^{-1}}{\sim} \Lambda^{\nu-1} R. \tag{1.31}$$

That means, for large R typical universes have the shape of long thin tubes. On the other hand, for $R \sim \Lambda^{-\nu}$ the exponential decay turns over into some power law and we get

$$\langle V \rangle_R \sim R^{\frac{1}{\nu}}, \text{ for } R \sim \Lambda^{-\nu}, \tag{1.32}$$

by inserting into (1.31). Per definition, the exponent of R in this equation equals the (grand canonical) intrinsic Hausdorff dimension $d_H = \frac{1}{\nu}$.

The large R behaviour of $G(V, R)$ can be computed from (1.30) by a saddle-point calculation. The result up to power corrections is:

$$G(V, R) \sim e^{-\tilde{c}\left(\frac{R}{V^\nu}\right)^{\frac{1}{1-\nu}}}, \text{ for } \frac{R}{V^\nu} \gg 1, \tag{1.33}$$

where $\tilde{c} = \frac{1-\nu}{\nu}(\nu c)^{\frac{1}{1-\nu}}$.

Another concept of fractal dimension can be applied in the ensemble of surfaces with fixed volume V for small distances R. Let us define with

$$l(g, R) = \frac{1}{V_g} \int_M d^2\xi \sqrt{g(\xi)} \int_M d^2\xi' \sqrt{g(\xi')}\, \delta(R - d_g(\xi, \xi')) \tag{1.34}$$

the average length of boundaries on a manifold M with the metric $g_{\mu\nu}$, which have a geodesic distance R from a marked point. Then $l(g, R)dR$ is the average volume of a spherical shell of radius R and thickness dR. The quantum expectation value in the ensemble of universes with fixed volume is given by

$$\langle l(g, R) \rangle_V = \frac{1}{Z(V)} \int \mathcal{D}[g_{\mu\nu}]_V\, l(g, R) = \frac{G(V, R)}{V Z(V)}. \tag{1.35}$$

Now we conclude from

$$\int_0^\infty dR\ \langle l(g, R) \rangle_V = V \tag{1.36}$$

that

$$\dim[V] = \dim[R] + \dim[\langle l(g, R) \rangle_V]. \tag{1.37}$$

Thus $\langle l(g,R)\rangle_V$ has the scaling behaviour

$$\langle l(g,R)\rangle_V = V^{1-\nu}\, F\left(\frac{R}{V^\nu}\right), \tag{1.38}$$

where $F(x)$ is a function which falls off exponentially for large x, see (1.33). The (canonical) intrinsic Hausdorff dimension d_h is now defined by the scaling of $\langle l(g,R)\rangle_V \sim R^{d_h-1}$ for small R. To be precise we expand (1.38) around $R=0$ and get:

$$\langle l(g,R)\rangle_V \sim V^{1-\nu d_h}\; R^{d_h-1}, \text{ for } R \ll V^{\frac{1}{d_h}}. \tag{1.39}$$

For smooth d-dimensional manifolds we have $d_H = d_h = d$. If $\langle l(g,R)\rangle_V$ stays nonzero and finite for $V \to \infty$ we have

$$\nu d_h = 1, \text{ that means } d_H = d_h. \tag{1.40}$$

This requirement is called the smooth fractal condition. It means that the average circumference of circles with a small geodesic radius R does not depend on the global volume of the universe. However it is well known that the smooth fractal condition is not fulfilled for the model of multicritical branched polymers [21], compare section 2.1.4. Therefore its validity is non-trivial and should not be taken for granted uncritically. It turns out that in two-dimensional quantum gravity the smooth fractal condition is fulfilled. That means that there is only one fractal dimension for short and for long distances together.

For pure gravity, in the case of spherical topology, the two-point function $G(\Lambda,R)$ can be calculated exactly with a transfer-matrix method [22] or alternatively by a peeling method [23]. The result

$$G(\Lambda,R) \propto \Lambda^{3/4}\frac{\cosh c\Lambda^{1/4}R}{\sinh^3 c\Lambda^{1/4}R} \tag{1.41}$$

fulfills the standard scaling relations, but with non-standard exponents. $G(\Lambda,R)$ falls off as $e^{-2c\Lambda^{1/4}R}$, from which we read off $\nu = 1/4$, so that the Hausdorff dimension of pure two-dimensional quantum gravity equals *four*.

Another scaling exponent, though not independent of γ and ν, can be defined by:

$$G(\Lambda,R) \sim R^{1-\eta}, \text{ for } 1 \ll R \ll \Lambda^{-\nu}. \tag{1.42}$$

η is called the anomalous scaling dimension. By expanding (1.41) for small R we get

$$G(\Lambda,R) = \frac{1}{R^3} - \frac{1}{15}\Lambda R + O(R^3), \tag{1.43}$$

and thus $\eta = 4$. This is a notable result, since in ordinary statistical systems η is always smaller than 2.

The so-called Fisher scaling relation

$$\gamma = \nu(2 - \eta) \tag{1.44}$$

relates the exponents defined above. It can be derived by applying dimensional arguments to (1.24).

1.2 Liouville theory: A brief reminder

A central problem in the continuum approach to two-dimensional quantum gravity is posed by the diffeomorphism invariance of the theory. It is comparatively easy [24] to derive an expression for the formal integration $\mathcal{D}g_{\mu\nu}$ since one can define a natural scalar product on the cotangent space to the space of all metrics which defines a volume form in the same way as in finite dimensional Riemannian geometry.

However, since the measure and the action are diffeomorphism invariant, these expressions are ill-defined. Therefore a gauge fixing and the factorization of the diffeomorphisms from the measure is required. This has been performed in [6] where the functional integration over geometries is expressed as a Liouville field theory. This theory has been developed and explained in [25, 26]. The measure for the Liouville mode is very complicated. Therefore two-dimensional quantum gravity has strictly speaking not been solved yet in the continuum approach.

However, the critical scaling exponents could be obtained in the light-cone gauge [7] and later in the conformal gauge by consistent scaling assumptions [8, 9].

1.2.1 Functional measures

Let $R(M)$ be the space of all positive definite Riemannian metrics on a d-dimensional manifold M and let $T_g R(M)$ be its cotangent space at a point $g_{\mu\nu} \in R(M)$.

To define the functional measure $\mathcal{D}g_{\mu\nu}$ one makes use of the fact, that the cotangent space $T_g R(M)$ can be naturally equipped with a diffeomorphism invariant scalar product $\langle \cdot, \cdot \rangle_T$. This is used to define a measure $\mathcal{D}_g \delta g_{\mu\nu}$ on $T_g R(M)$ which descends to a measure on $R(M)$. The line element ds^2 for

fluctuations $\delta g_{\mu\nu} \in T_g R(M)$ of the metric can be written as

$$ds^2 = \langle \delta g, \delta g \rangle_T \equiv \int_M d^d\xi \sqrt{g}\ \delta g_{\mu\nu} G^{\mu\nu,\alpha\beta} \delta g_{\alpha\beta}, \tag{1.45}$$

with the DeWitt (super)metric [27]

$$G^{\mu\nu,\alpha\beta} = \frac{1}{2}(g^{\mu\alpha}g^{\nu\beta} + g^{\mu\beta}g^{\nu\alpha} + C g^{\mu\nu}g^{\alpha\beta}). \tag{1.46}$$

Up to an overall normalization constant, (1.45) is the only diffeomorphism invariant, ultralocal, that means dynamically neutral distance on $T_g R(M)$. The constant C in (1.46) takes the value -2 in canonical quantum gravity [27]. In our framework C cannot be computed. The choice of C determines the signature of the metric $\langle \cdot, \cdot \rangle_T$ as can be seen by splitting the fluctuation $\delta g_{\mu\nu}$ into its trace part $g_{\mu\nu}\delta c$ and its tracefree part $\delta h_{\mu\nu}$. This decomposition is orthogonal with respect to $\langle \cdot, \cdot \rangle_T$. On the tracefree subspace $G^{\mu\nu,\alpha\beta}$ has eigenvalue 1, while it has eigenvalue $1 + \frac{Cd}{2}$ on the trace sector. Thus, the DeWitt metric is positive definite for $C > -\frac{2}{d}$ and negative definite for $C < -\frac{2}{d}$. For $C = -\frac{2}{d}$, $G^{\mu\nu,\alpha\beta}$ is the orthogonal projection on the tracefree part of $\delta g_{\mu\nu}$. The inverse of the DeWitt metric, which satisfies

$$G_{\mu\nu,\alpha\beta} G^{\alpha\beta,\rho\sigma} = \frac{1}{2}\left(\delta^\rho_\mu \delta^\sigma_\nu + \delta^\sigma_\mu \delta^\rho_\nu\right), \tag{1.47}$$

is given by

$$G_{\mu\nu,\alpha\beta} = \frac{1}{2}\left(g_{\mu\alpha}g_{\nu\beta} + g_{\mu\beta}g_{\nu\alpha} - \frac{C}{1 + \frac{Cd}{2}} g_{\mu\nu}g_{\alpha\beta} \right), \tag{1.48}$$

as can be verified by multiplication with (1.46).

The line element (1.45) can be rewritten as

$$ds^2 = \int_M d^d\xi \int_M d^d\xi' \sum_{\mu\le\nu} \sum_{\alpha\le\beta} \delta g_{\mu\nu}(\xi) \Omega^{\mu\nu,\alpha\beta}(\xi,\xi') \delta g_{\alpha\beta}(\xi'), \tag{1.49}$$

with $\Omega^{\mu\nu,\alpha\beta}(\xi,\xi') = \sqrt{g(\xi)}\, \Gamma^{\mu\nu,\alpha\beta}(\xi)\ \delta(\xi - \xi')$, and the $\frac{d(d+1)}{2} \times \frac{d(d+1)}{2}$ matrix $\Gamma^{\mu\nu,\alpha\beta}(\xi)$ is defined as

$$\Gamma^{\mu\nu,\alpha\beta}(\xi) = \begin{cases} 4G^{\mu\nu,\alpha\beta}(\xi), & \text{if } \mu < \nu \text{ and } \alpha < \beta, \\ 2G^{\mu\nu,\alpha\beta}(\xi), & \text{if } \mu < \nu \text{ and } \alpha = \beta, \text{ or } \mu = \nu \text{ and } \alpha < \beta, \\ G^{\mu\nu,\alpha\beta}(\xi), & \text{if } \mu = \nu \text{ and } \alpha = \beta. \end{cases} \tag{1.50}$$

This induces a volume form $\mathcal{D}\delta g_{\mu\nu}$ on the cotangent space, which descends to a volume form $\mathcal{D}g_{\mu\nu}$ on $R(M)$:

$$\mathcal{D}\delta g_{\mu\nu} = \sqrt{\det \Omega^{\mu\nu,\alpha\beta}(\xi,\xi')} \prod_{\xi\in M,\ \mu\le\nu} d\delta g_{\mu\nu}(\xi), \tag{1.51}$$

$$\mathcal{D}g_{\mu\nu} = \sqrt{\det \Omega^{\mu\nu,\alpha\beta}(\xi,\xi')} \prod_{\xi\in M,\ \mu\le\nu} dg_{\mu\nu}(\xi). \tag{1.52}$$

We first note that due to the δ-function the determinant of $\Omega^{\mu\nu,\alpha\beta}(\xi,\xi')$ factorizes into a product over all spacetime points $\xi \in M$. The local determinant of the $\frac{d(d+1)}{2} \times \frac{d(d+1)}{2}$ matrix $\sqrt{g(\xi)}\ \Gamma^{\mu\nu,\alpha\beta}(\xi)$ can be evaluated by variation with respect to $\delta \log g = -g^{\mu\nu}\delta g_{\mu\nu}$:

$$\begin{aligned}
\delta \log \det\left(\sqrt{g}\ \Gamma^{\mu\nu,\alpha\beta}\right) &= \mathrm{Tr}\ \delta \log\left(\Gamma^{\mu\nu,\alpha\beta}\right) + \mathrm{Tr}\ \delta \log\left(\sqrt{g}\ 1^{\mu\nu,\alpha\beta}\right) \\
&= G_{\mu\nu,\alpha\beta}\delta G^{\alpha\beta,\mu\nu} + \frac{d(d+1)}{4}\delta \log g \\
&= \frac{1}{4}(d+1)(d-4)\ \delta \log g,
\end{aligned} \tag{1.53}$$

where it has been used that the inverse of $\Gamma^{\mu\nu,\alpha\beta}$ equals $G_{\mu\nu,\alpha\beta}$ restricted to all indices with $\mu \le \nu$ and $\alpha \le \beta$. Therefore it follows that

$$\det\left(\sqrt{g}\ \Gamma^{\mu\nu,\alpha\beta}\right) = \kappa g^{\frac{(d+1)(d-4)}{4}}, \tag{1.54}$$

where the constant κ can be determined by specializing to $g_{\mu\nu} = \delta_{\mu\nu}$, which gives $\kappa = \left(1 + \frac{Cd}{2}\right)/2^{\frac{d(d-1)}{2}}$. Thus the measures (1.51) and (1.52) equal

$$\mathcal{D}\delta g_{\mu\nu} = \mathrm{const} \times \prod_{\xi\in M}\prod_{\mu\le\nu} g(\xi)^{\frac{d-4}{4d}} d\delta g_{\mu\nu}(\xi), \tag{1.55}$$

$$\mathcal{D}g_{\mu\nu} = \mathrm{const} \times \prod_{\xi\in M}\prod_{\mu\le\nu} g(\xi)^{\frac{d-4}{4d}} dg_{\mu\nu}(\xi), \tag{1.56}$$

which we normalize such that

$$\int \mathcal{D}\delta g_{\mu\nu}\ e^{-\frac{1}{2}\langle \delta g, \delta g\rangle_T} = 1. \tag{1.57}$$

Note that $\sigma = \frac{d-4}{4d}$ is the only exponent of $g(\xi)$ in (1.55) or (1.56) such that these measures are diffeomorphism invariant. With the measure

$$\prod_{\xi\in M}\prod_{\mu\le\nu} g(\xi)^{\sigma} d\delta g_{\mu\nu}(\xi), \tag{1.58}$$

the Gaussian integral (1.57) is resolved by computing the determinant of $g(\xi)^{\frac{1}{2}-2\sigma}\Gamma^{\mu\nu,\alpha\beta}$. This can be done along the lines of (1.53). The result is

$$\det\left(g(\xi)^{\frac{1}{2}-2\sigma}\Gamma^{\mu\nu,\alpha\beta}\right) = \mathrm{const} \times g(\xi)^{\frac{(d+1)((1-4\sigma)d-4)}{4}}, \tag{1.59}$$

which is a diffeomorphism invariant constant if and only if $\sigma = \frac{d-4}{4d}$. The same result can be derived more rigorously by using the BRS-symmetry associated with general coordinate transformations [28, 29].

Analogously, the measures for scalar and vector fields can be derived. The corresponding metrics $\langle\cdot,\cdot\rangle_S$ and $\langle\cdot,\cdot\rangle_V$ are defined as

$$\langle\phi,\phi\rangle_S = \int_M d^d\xi\sqrt{g}\ \phi(\xi)\phi(\xi), \tag{1.60}$$

$$\langle\psi,\psi\rangle_V = \int_M d^d\xi\sqrt{g}\ \psi_\mu(\xi)g^{\mu\nu}\psi_\nu(\xi). \tag{1.61}$$

The measures $\mathcal{D}_g\phi$ and $\mathcal{D}_g\psi_\mu$ turn out as:

$$\mathcal{D}_g\phi = \prod_{\xi\in M} g(\xi)^{\frac{1}{4}}d\phi(\xi), \tag{1.62}$$

$$\mathcal{D}_g\psi_\mu = \prod_{\xi\in M}\prod_{\mu=1}^{d} g(\xi)^{\frac{d-2}{4d}}d\psi_\mu(\xi). \tag{1.63}$$

1.2.2 Factorization of the diffeomorphisms

By construction, the functional measure and the action in (1.1) and (1.10) are invariant under diffeomorphisms. Therefore, the measure overcounts physically equivalent configurations related by the group of diffeomorphisms. To avoid this overcounting of gauge equivalent metrics, the diffeomorphisms have to be factored from the measure. In two dimensions, this can be done in a covariant way in the conformal gauge.

Let us first recall that the space of metrics on M modulo diffeomorphisms and Weyl transformations is a finite dimensional compact space, called the moduli space $\mathcal{M}$ of M, which has dimension 0, 2 and $6h-6$ for genus 0, 1 and $h \geq 2$ respectively. It is parametrized by the Teichmüller parameters $\tau = (\tau_1,\ldots,\tau_N)$. That means that if for each moduli $\tau \in \mathcal{M}$ we choose a fixed background metric $\hat{g}_{\mu\nu}(\tau)$, all other metrics $g_{\mu\nu}$ are contained in the orbits under diffeomorphisms and Weyl transformations:

$$g_{\mu\nu} = f^\star\left(e^\phi\hat{g}_{\mu\nu}(\tau)\right), \tag{1.64}$$

where $f^\star$ denotes the action of the diffeomorphism $f : M \to M$. (Actually the background metrics $\hat{g}_{\mu\nu}$ can be chosen such that they all have constant curvature.)

Any infinitesimal change $\delta g_{\mu\nu}$ in the metric can be decomposed into an infinitesimal Weyl transformation $\delta\phi$, the effect of an infinitesimal diffeo-

morphism ξ, and the effect of varying the Teichmüller parameters:

$$\delta g_{\mu\nu} = f^\star \delta\phi \; g_{\mu\nu} + (\nabla_\mu \xi_\nu + \nabla_\nu \xi_\mu) + \frac{\partial g_{\mu\nu}}{\partial \tau_i} \delta\tau_i. \tag{1.65}$$

Here ∇_μ denotes the covariant derivative. We want to orthogonalize this decomposition with respect to the scalar product $\langle \cdot, \cdot \rangle_T$. First note that the tracefree part of the effect of the diffeomorphism ξ is given by the conformal Killing form

$$(P\xi)_{\mu\nu} = \nabla_\mu \xi_\nu + \nabla_\nu \xi_\mu - g_{\mu\nu} \nabla_\alpha \xi^\alpha, \tag{1.66}$$

which maps vector fields into traceless symmetric tensors. The zero modes of P are called conformal Killing vectors. For spherical topology there are six linearly independent conformal Killing vectors, for torus topology there are two while there are none for higher genus. The conformal Killing vectors are important because they induce a variation $\delta g_{\mu\nu}$ in the direction of $g_{\mu\nu}$ which can thus be compensated by a Weyl rescaling. In other words, the decomposition (1.65) is in general not unique. To make it unique one chooses a diffeomorphism $\tilde{\xi} \in (\ker P)^\perp$ orthogonal to the zero modes of P. That means the gauge will only be fixed up to the conformal Killing vectors and we expect a remaining, at most six-dimensional symmetry in the gauge-fixed expressions. It follows that all variations $\delta g_{\mu\nu} \in T_g R(M)$ of the metric can be uniquely decomposed into the variation of the conformal factor, the action of a diffeomorphism orthogonal to the conformal Killing vectors and the variation of the Teichmüller parameters. This decomposition can be written as a mutually orthogonal sum of a trace part, a tracefree part orthogonal to the moduli deformations and these moduli deformations:

$$\begin{aligned} \delta g_{\mu\nu} \;=\; & \{\tilde{f}^\star \delta\phi + \nabla_\alpha \tilde{\xi}^\alpha + \frac{1}{2} g^{\alpha\beta} \frac{\partial g_{\alpha\beta}}{\partial \tau_i} \delta\tau_i\} g_{\mu\nu} + \{P\tilde{\xi} + P(P^+P)^{-1} P^+ k^i_{\mu\nu} \delta\tau_i\} \\ & + \left(1 - P(P^+P)^{-1} P^+\right) k^i_{\mu\nu} \delta\tau_i, \end{aligned} \tag{1.67}$$

where

$$k^i_{\mu\nu} = \frac{\partial g_{\mu\nu}}{\partial \tau_i} - \frac{1}{2} g_{\mu\nu} g^{\alpha\beta} \frac{\partial g_{\alpha\beta}}{\partial \tau_i}. \tag{1.68}$$

The adjoint P^+ of P is defined by the relation $\langle h, P\xi \rangle_T = \langle P^+ h, \xi \rangle_V$. To prove that the decomposition (1.67) is orthogonal just note that $1 - P(P^+P)^{-1}P^+$ is the projector on the zero modes ψ_l of P^+. The tracefree part is orthogonal to the trace part by definition of the scalar product $\langle \cdot, \cdot \rangle_T$.

The change of variables $g_{\mu\nu} \to (\phi, \tilde{f}, \tau)$ involves a Jacobian $J(\phi, \tau)$:

$$\int \mathcal{D}[g_{\mu\nu}]\ \mathcal{F}(g) = \int \mathcal{D}f \int \frac{\mathcal{D}\tilde{f}}{\mathcal{D}f} \mathcal{D}_{e^{\phi}\hat{g}}\phi d\tau\ J(\phi,\tau)\ \mathcal{F}(e^{\phi}\hat{g}(\tau)). \tag{1.69}$$

Here one integrates some reparametrization invariant functional $\mathcal{F}$. To find J one substitutes the orthogonal decomposition (1.67) into the normalization condition

$$1 = \int \mathcal{D}\delta g_{\mu\nu}\ e^{-\frac{1}{2}\langle \delta g, \delta g\rangle_T} \tag{1.70}$$

and performs the change of variables on the tangent space. The result of these Gaussian integrations is

$$1 = J(\phi,\tau)\left[\frac{\det \langle \psi_k, \psi_l\rangle_T}{\det'(P^+P)}\right]^{\frac{1}{2}} \left[\det \left\langle \psi_m, \frac{\partial g}{\partial \tau_n}\right\rangle_T\right]^{-1}. \tag{1.71}$$

At last, the change from the diffeomorphisms f to $\tilde{f}$ and the conformal Killing vectors ω_a, which fulfill $P(e^{\phi}\omega_a) = 0$, has to be computed. The final result is

$$\int \mathcal{D}[g_{\mu\nu}]\ \mathcal{F}(g) = \int \mathcal{D}f_{\mu} \int \frac{d\tau}{\nu(\tau)} \det \left\langle \psi_m, \frac{\partial g}{\partial \tau_n}\right\rangle_T \times \int \mathcal{D}_{e^{\phi}\hat{g}}\phi \left[\frac{\det'(P^+P)}{\det \langle \psi_k, \psi_l\rangle_T \det \langle \omega_a, \omega_b\rangle_V}\right]^{\frac{1}{2}} \mathcal{F}(e^{\phi}\hat{g}(\tau)), \tag{1.72}$$

where $\nu(\tau)$ is the volume of the group generated by the conformal Killing vectors. Now the diffeomorphisms can be factored out. Since it turns out that $\frac{d\tau}{\nu(\tau)} \det \left\langle \psi_m, \frac{\partial g}{\partial \tau_n}\right\rangle_T$ depends only on the Teichmüller parameters and not on ϕ we denote this as $\mathcal{D}\tau$. The square root is called the Faddeev-Popov determinant and denoted as $Z_{FP}(e^{\phi}\hat{g})$.

1.2.3 Gravitational dressing of scaling exponents

After factoring the diffeomorphisms from the measure the gauge-fixed expression for the partition function (1.10) is:

$$Z(V) = \int \mathcal{D}\tau \int \mathcal{D}_{e^{\phi}\hat{g}}\phi\ \delta(V - V_{e^{\phi}\hat{g}})\ Z_{FP}(e^{\phi}\hat{g}) Z_{mat}(e^{\phi}\hat{g}), \tag{1.73}$$

where Z_{mat} is defined as:

$$Z_{mat}(e^{\phi}\hat{g}) = \int \mathcal{D}_{e^{\phi}\hat{g}}X\ e^{-S_{matter}(e^{\phi}\hat{g}, X)}, \tag{1.74}$$

with some matter action for matter with central charge D. The measure for the Liouville mode ϕ is defined by the line element

$$d^2s = \langle \delta\phi, \delta\phi \rangle_S = \int_M d^2\xi \sqrt{\hat{g}}\ e^{\phi}\ \delta\phi(\xi)\delta\phi(\xi), \tag{1.75}$$

which depends on ϕ itself. Therefore the corresponding measure $\mathcal{D}_{e^{\phi}\hat{g}}\phi$ is very complicated and it is unknown how to perform the functional integration over ϕ with this measure. The idea used to overcome these problems is to go over to the translational invariant measure $\mathcal{D}_{\hat{g}}\phi$ and to shift all dependence on ϕ in (1.73) into the action.

For this transformation we use the relations

$$Z_{mat}(e^{\phi}\hat{g}) = Z_{mat}(\hat{g}) e^{\frac{D}{48\pi} S_L(\hat{g},\phi)}, \tag{1.76}$$

$$Z_{FP}(e^{\phi}\hat{g}) = Z_{FP}(\hat{g}) e^{-\frac{26}{48\pi} S_L(\hat{g},\phi)}, \tag{1.77}$$

$$\mathcal{D}_{e^{\phi}\hat{g}}\phi = \mathcal{D}_{\hat{g}}\phi\ e^{\frac{1}{48\pi} S_L(\hat{g},\phi)}, \tag{1.78}$$

where $S_L(\hat{g},\phi)$ is the Liouville action:

$$S_L(\hat{g},\phi) = \int_M d^2\xi \sqrt{\hat{g}}\ \left(\frac{1}{2} \hat{g}^{\mu\nu} \partial_\mu\phi\partial_\nu\phi + \hat{\mathcal{R}}\phi \right). \tag{1.79}$$

Here $\hat{\mathcal{R}}$ is the curvature scalar in the metric $\hat{g}_{\mu\nu}$.

The relation (1.76) for the matter partition function can be derived by a couple of methods such as the heat kernel regularization [30] or the zeta function regularization [31]. Since the matter action is invariant under Weyl transformations we can actually say that the whole contribution comes from the transformation of the measure $\mathcal{D}_{e^{\phi}\hat{g}}X$ under conformal rescalings of the metric. This is also the reason for the trace anomaly of the energy momentum tensor.

The transformation property (1.77) for the Faddeev-Popov determinant can be derived in a similar way. However, the involved operators are much more complicated, compare (1.72).

The transformation (1.78) could not be derived rigorously by these or other methods. David [8] and Distler and Kawai [9] use a selfconsistent bootstrap method to find this relation. Under the assumption that the factor takes the form of an exponential of a Liouville action with arbitrary coefficients the coefficients are determined from general invariance considerations.

In this calculation the Liouville mode ϕ is rescaled by a factor α, which has to be determined. Taking all factors together, the partition function

(1.73) can be rewritten as:

$$Z(V) = \int \mathcal{D}\tau \int \mathcal{D}_{\hat{g}}\phi \; Z_{FP}(\hat{g}) \; Z_{mat}(\hat{g}) \; e^{\frac{D-25}{48\pi} S_L(\hat{g},\phi)} \; \delta(V - V_{e^{\alpha\phi}\hat{g}}), \tag{1.80}$$

where the measures depend only on the background metric $\hat{g}$. Thus the partition function scales under a rescaling of the volume as:

$$Z(\lambda V) = \lambda^{\frac{D-25}{12\alpha}\chi - 1} Z(V), \tag{1.81}$$

where χ is the Euler characteristic of the manifold M. To find (1.81) we have shifted ϕ by $\frac{1}{\alpha}\log\lambda$ and used the Gauss-Bonnet theorem (1.3).

Now we have to compute the dressing exponent α. It turns out that more generally we can compute the anomalous scaling dimensions for a whole family of operators. Consider the observable

$$\mathcal{O}(g) = \int_M d^2\xi \sqrt{g} \; \Phi(g), \tag{1.82}$$

where Φ is a spinless primary field of dimension Δ_0. Typical simple examples are the volume $V_g = \int_M d^2\xi\sqrt{g}$ with $\Delta_0 = 0$ and the identity $1 = \int_M d^2\xi\sqrt{g} \; \frac{1}{\sqrt{g}}\delta(\xi)$ with $\Delta_0 = 1$. Then $\mathcal{O}(g)$ transforms under a rescaling $g_{\mu\nu} \to \lambda g_{\mu\nu}$ of the metric as

$$\mathcal{O}(\lambda g) = \lambda^{1-\Delta_0}\mathcal{O}(g), \tag{1.83}$$

if the geometry is kept fixed. Under the average over all geometries the scaling behaviour exponent Δ_0 is in general changed to some other value Δ:

$$\langle \mathcal{O}(g) \rangle_{\lambda V} = \lambda^{1-\Delta} \langle \mathcal{O}(g) \rangle_V . \tag{1.84}$$

The coupling to gravity dresses the scaling behaviour of the field Φ. Therefore we make the ansatz

$$\mathcal{O}(\hat{g}, \phi) = \int_M d^2\xi \sqrt{g} \; e^{\beta\phi} \Phi(\hat{g}) \tag{1.85}$$

for the observable after the transition to the translational invariant measure. Note that $\mathcal{O}(\hat{g}, \phi)$ is diffeomorphism invariant and fulfills $\mathcal{O}(\hat{g}, 0) = \mathcal{O}(\hat{g})$. Now observe that (1.76), (1.77), (1.78) and $\mathcal{O}(e^{\phi}\hat{g})$ are invariant under the transformation

$$\hat{g}_{\mu\nu}(\xi) \to e^{\sigma(\xi)}\hat{g}_{\mu\nu}(\xi), \quad \phi(\xi) \to \phi(\xi) - \sigma(\xi). \tag{1.86}$$

Therefore also $\mathcal{O}(\hat{g},\phi)$ has to be invariant. This requirement leads to an expression for the gravitational dressing exponent β as follows. Under (1.86) the product $\sqrt{\hat{g}}\ \Phi(\hat{g})$ transforms as

$$\sqrt{\hat{g}(\xi)}\Phi(\hat{g}) \to e^{(1-\Delta_0)\sigma(\xi)}\sqrt{\hat{g}(\xi)}\Phi(\hat{g}). \tag{1.87}$$

Thus $e^{\beta\phi(\xi)}$ has to transform as

$$e^{\beta\phi(\xi)} \to e^{(\Delta_0-1)\sigma(\xi)}e^{\beta\phi(\xi)}. \tag{1.88}$$

On the other hand we can determine the transformation of $e^{\beta\phi(\xi)}$ by computing the Gaussian integral

$$F_\phi(\hat{g},\beta) = \int \mathcal{D}_{\hat{g}}\phi\ e^{\frac{D-25}{48\pi}S_L(\hat{g},\phi)+\beta\phi(\xi)}, \tag{1.89}$$

where $\beta\phi(\xi)$ is interpreted as an additional δ-function source. If we define

$$\hat{J}(x) = \hat{\mathcal{R}}(x) + \frac{48\pi\beta}{D-25}\frac{1}{\sqrt{\hat{g}(x)}}\delta(x-\xi), \tag{1.90}$$

we get:

$$\begin{aligned} F_\phi(\hat{g},\beta) &= \exp\left[\frac{D-25}{96\pi}\int_M d^2x\sqrt{g}\ \hat{J}(x)(\Delta_{\hat{g}}^{-1}\hat{J})(x)\right] \qquad (1.91)\\ &= F_\phi(\hat{g},0)\ \exp\left[\beta\int_M d^2x\sqrt{\hat{g}}\ \Delta_{\hat{g}}^{-1}(\xi,x)\hat{\mathcal{R}}(x)\ +\ \frac{24\pi\beta^2}{D-25}\Delta_{\hat{g}}^{-1}(\xi,\xi)\right]. \end{aligned}$$

Under the transformation (1.86) the first term in the exponent gives an extra $-\beta\sigma(\xi)$. The second term is an evaluation of the propagator at coinciding points which naively seen is singular. Thus we have to renormalize this expression [32]. The renormalized propagator is defined as

$$\Delta_{\hat{g},R}^{-1}(\xi,\xi) = \lim_{x\to\xi}\left(\Delta_{\hat{g}}^{-1}(x,\xi) - \frac{1}{4\pi}\log d_{\hat{g}}(x,\xi)^2\right), \tag{1.92}$$

where $d_{\hat{g}}(x,\xi)$ is the geodesic distance between x and ξ. Therefore the second term in the exponent gains an extra $\frac{6\beta^2}{25-D}\sigma(\xi)$ under (1.86). Taking both contributions together we conclude:

$$e^{\beta\phi(x)} \to e^{-\beta\sigma(x)+\frac{6\beta^2}{25-c}\sigma(x)}\ e^{\beta\phi(x)} \stackrel{!}{=} e^{(\Delta_0-1)\sigma(x)}\ e^{\beta\phi(x)}. \tag{1.93}$$

The solution to this quadratic equation determines β and thus also α:

$$\begin{aligned} \beta &= \frac{1}{12}\left(25-D-\sqrt{(25-D)(1-D+24\Delta_0)}\right), \qquad (1.94)\\ \alpha &= \frac{1}{12}\left(25-D-\sqrt{(25-D)(1-D)}\right). \qquad (1.95) \end{aligned}$$

The sign of the square root is determined by requesting that in the classical limit $c \to -\infty$ the scaling of observables is the same as in a fixed geometry. With α inserted into (1.81) we can now verify the form (1.14) for the susceptibility exponent γ.

For the expectation values of observables the dependence on the topology cancels , and we get:

$$\langle \mathcal{O}(g) \rangle_{\lambda V} = \lambda^{\frac{\beta}{\alpha}} \langle \mathcal{O}(g) \rangle_V = \lambda^{1-\Delta} \langle \mathcal{O}(g) \rangle_V . \tag{1.96}$$

Δ evaluates as

$$\Delta = \frac{\sqrt{1 - D + 24\Delta_0} - \sqrt{1 - D}}{\sqrt{25 - D} - \sqrt{1 - D}}. \tag{1.97}$$

The expectation value of products of observables scales with the sum of their scaling exponents.

1.3 Fractal dimensions

Quantum gravity as defined above is a functional integration over geometries. Therefore we have to ask which geometrical concepts survive the quantum average and how geometry is changed by it. In any theory of quantum gravity the structure of spacetime is of primary interest and even its dimensionality is a *dynamical* quantity. The transfer matrix method in [22] applied to pure two-dimensional quantum gravity revealed that the spacetime has a self-similar structure on all scales with a fractal dimension *four*. An understanding of the quantum spacetime in two-dimensional quantum gravity might help to understand similar dynamical changes of the geometry in higher dimensions.

In this section we introduce three different concepts of dimensionality to characterize the fractal structure of spacetime, each of which might describe different aspects of the geometry. The intrinsic Hausdorff dimension or fractal dimension d_H has already been defined above by the scaling of observables such as the average circumference of circles with their "geodesic" radii. These measurements are performed within the intrinsic geometry of the universes. They are analogous to the usual "clock and rod" measurements in classical gravity. General results for d_H from Liouville theory have been given in [33].

We can also characterize the fractal structure of spacetime by embedding it into a D-dimensional, typically Euclidean flat space. Measurements of, for example, the mean square extent of the universes are then performed

with the metric of the embedding space. The finite size scaling of the mean square extent of the surfaces defines the extrinsic Hausdorff dimension D_H. Intuitively the embedding gives us an idea what typical universes would "look like". Indeed it is known that the extrinsic Hausdorff dimension of two-dimensional quantum gravity is infinity [34, 35]. That means that the surfaces are crumpled together in a complicated way.

Instead of determining the metrical structure of the spacetime directly by putting out clocks and rods, one can also choose to observe the propagation of test particles. From such observations one can in the classical case determine the propagator and via an inverse Laplace transformation the associated heat kernel. Since all this can be done in a reparametrization invariant way, we can define corresponding quantum observables. This leads to the definition of the spectral dimension d_s as the scaling exponent of the short time return probability distribution of diffusing matter. The first general analytical result for the spectral dimension of two-dimensional quantum gravity coupled to conformal matter with central charge D was given in [1].

1.3.1 Extrinsic Hausdorff dimension

The extrinsic Hausdorff dimension D_H of a fractal object embedded in a D-dimensional space is a measure for the extent of the object measured with the metric of the embedding space. D_H can be defined for mathematical fractals as well as for random surfaces and for quantum gravity. Let us focus on quantum gravity. The partition function for two-dimensional quantum gravity with fixed volume V of spacetime coupled to D Gaussian matter fields is

$$Z(V) = \int \mathcal{D}[g_{\mu\nu}]_V \int \mathcal{D}_g X_\mu \big|_{cm} e^{-\int_M d^2\xi \sqrt{g}\ g^{\alpha\beta}\partial_\alpha X_\mu \partial_\beta X_\mu}. \tag{1.98}$$

Here $\mathcal{D}_g X_\mu\big|_{cm}$ denotes the functional integration over the D matter fields such that the centre of mass is fixed to zero, which amounts in dropping the zero modes of the Laplacian when integrating over the matter fields. This is a slight variation of (1.10) which allows us to define the extrinsic Hausdorff dimension as

$$\langle X^2 \rangle_V \sim V^{\frac{2}{D_H}}, \text{ for } V \to \infty, \tag{1.99}$$

with the definition

$$\begin{aligned} \langle X^2 \rangle_V \ \equiv\ & \frac{1}{Z(V)} \int \mathcal{D}[g_{\mu\nu}]_V \int \mathcal{D}_g X_\mu \big|_{cm} e^{-\int_M d^2\xi \sqrt{g}\ g^{\alpha\beta}\partial_\alpha X_\mu \partial_\beta X_\mu} \\ & \times \frac{1}{DV} \int_M d^2\xi \sqrt{g}\ X_\mu^2(\xi), \end{aligned} \tag{1.100}$$

which is equivalent to

$$\begin{aligned}\left\langle X^2\right\rangle_V &\equiv \frac{1}{Z(V)}\int \mathcal{D}[g_{\mu\nu}]_V \int \mathcal{D}_g X_\mu\big|_{cm} e^{-\int_M d^2\xi\sqrt{g}\ g^{\alpha\beta}\partial_\alpha X_\mu \partial_\beta X_\mu} \\ &\quad \times \frac{1}{2DV^2}\int_M d^2\xi_1\sqrt{g}\int_M d^2\xi_2\sqrt{g}\ (X(\xi_1)-X(\xi_2))^2. \quad (1.101)\end{aligned}$$

To understand the definition of D_H note that the fields X_μ define an embedding of the manifold M in a D-dimensional Euclidean space. That means that we measure the Euclidean mean-square extent of spacetime embedded in D dimensions. Mathematically more conservatively the extrinsic Hausdorff dimension is defined by constructing a covering of the geometrical object embedded in some D-dimensional space by a union of n small balls. n depends on the typical macroscopic scale r of the system and scales with the dimension D_H and is related to the mean square extent of the system as

$$n \sim r^{D_H} \sim \left\langle X^2\right\rangle^{\frac{D_H}{2}}. \quad (1.102)$$

The extrinsic Hausdorff dimension for two-dimensional quantum gravity has been computed in [34, 35]. Note that the treatment in [34] is based on a more general scaling assumption than is needed in two-dimensional quantum gravity. Their general result was specialized to quantum gravity in [35]. In the following we briefly summarize the derivation.

If we define the two-point function by

$$G(p) = \left\langle \int_M d^2\xi_1\sqrt{g}\int_M d^2\xi_2\sqrt{g}\ e^{ip(X(\xi_1)-X(\xi_2))}\right\rangle_V, \quad (1.103)$$

it follows that

$$\left\langle X^2\right\rangle_V = -\frac{1}{2DV^2}\frac{\partial^2}{\partial p^2}G(p)\big|_{p=0}. \quad (1.104)$$

In flat space the two-point function behaves as $V^{2-\Delta_0(p)}$ with $\Delta_0 \sim p^2$. Coupled to gravity this acquires a gravitational dressing which can be computed to be:

$$\Delta(p) = \frac{\sqrt{1-D+24\Delta_0(p)}-\sqrt{1-D}}{\sqrt{25-D}-\sqrt{1-D}}. \quad (1.105)$$

If we differentiate $G(p) \underset{V\to\infty}{\sim} V^{2-\Delta(p)}$ two times with respect to p at $p=0$ we get

$$\begin{aligned}\left\langle X^2\right\rangle_V &\sim \log V, \text{ for } D<1, \quad (1.106)\\ \left\langle X^2\right\rangle_V &\sim \log^2 V, \text{ for } D=1, \quad (1.107)\end{aligned}$$

for large volumes V. In both cases $\langle X^2 \rangle_V$ grows slower than any power of V. Thus we conclude that $D_H = \infty$. For spherical topology this has already been found for $D \to -\infty$ and to one-loop order in [36].

1.3.2 Spectral dimension

To probe the structure of classical spacetime one can study the properties of propagating particles, typically of free particles or diffusing matter. Let $\Psi(\xi, T)$ be the wave function for diffusion on the d-dimensional compact manifold M. Ψ depends on the time-parameter T and is a function of the points in M which fulfills the diffusion equation

$$\frac{\partial}{\partial T}\Psi(\xi, T) = \Delta_g \Psi(\xi, T) \tag{1.108}$$

for the initial condition

$$\Psi(\xi, 0) = \frac{1}{\sqrt{g(\xi)}}\delta(\xi_0 - \xi). \tag{1.109}$$

Here Δ_g is the Laplace-Beltrami operator corresponding to the metric $g_{\mu\nu}$ of M. The solution of the diffusion equation can be written as

$$\Psi(\xi, T) = e^{T\Delta'_g}\Psi(\xi, 0) = \int_M d^d\xi' \sqrt{g}\; K'_g(\xi, \xi'; T)\Psi(\xi', 0), \tag{1.110}$$

which defines the probability distribution (or heat kernel) $K'_g(\xi, \xi'; T)$ for diffusion on a compact manifold M with metric $g_{\mu\nu}$. Here and in the following we take into account, that the Laplace operator Δ_g on compact surfaces has zero modes which should be projected out. This is indicated with a prime. $K'_g(\xi, \xi'; T)$ is related to the massless scalar propagator $(-\Delta_g)^{-1}$ by

$$\langle \xi' | (-\Delta_g)^{-1} | \xi \rangle' = \int_0^\infty dT\; K'_g(\xi, \xi'; T). \tag{1.111}$$

It is known that the average return probability distribution

$$RP'_g(T) \equiv \frac{1}{V_g}\int_M d^d\xi \sqrt{g}\; K'_g(\xi, \xi; T) \tag{1.112}$$

at time T for diffusion on a d-dimensional manifold with a smooth geometry $[g_{\mu\nu}]$ admits the following asymptotic expansion for small times T:

$$RP'_g(T) \sim \frac{1}{T^{d/2}}\sum_{r=0}^{\infty} a_n T^n = \frac{1}{T^{d/2}}(1 + O(T)), \tag{1.113}$$

where the coefficients a_n are *diffeomorphism invariant* integrals of polynomials in the metric and the curvature and in their covariant derivatives, see for example [37]. This asymptotic expansion breaks down when T is of the order $V^{\frac{2}{d}}$, when the exponential decay of the heat kernel becomes dominant.

Since $RP'_g(T)$ is reparametrization invariant, we can define its quantum average over geometries for fixed volume V of spacetime as:

$$RP'_V(T) \equiv \frac{1}{Z(V)} \int \mathcal{D}[g_{\mu\nu}]_V \, e^{-S_{\text{eff}}(g)} RP'_g(T). \tag{1.114}$$

Here $S_{\text{eff}}(g)$ denotes the effective action of quantum gravity after integrating out possible matter fields.

The spectral dimension d_s is now defined by the short time asymptotic behaviour of $RP'_V(T)$:

$$RP'_V(T) \sim \frac{1}{T^{d_s/2}}(1 + O(T)). \tag{1.115}$$

It is natural to expect that under the average (1.114) over all geometries the only remaining geometric invariant is the fixed volume V. Thus we expect that $RP'_V(T)$ has the form

$$RP'_V(T) = \frac{1}{T^{d_s/2}} F\left(\frac{T}{V^{2/d_s}}\right), \tag{1.116}$$

where $F(0) > 0$ and $F(x)$ falls off exponentially for $x \to \infty$. This scaling assumption is the main input into our derivation of the spectral dimension of two-dimensional quantum gravity. It is very well documented in numerical simulations [38, 39].

For a fixed smooth geometry the spectral dimension is by definition equal to the dimension d of the manifold. However, the quantum average can a priori change this behaviour. Actually we know that the intrinsic Hausdorff dimension of two-dimensional quantum gravity is different from two. In this sense generic geometries are fractal with probability one. For diffusion on fixed (often embedded) fractal structures it is well known, that the spectral dimension can be different from the embedding dimension as well as from the fractal dimension. The diffusion law, measured in the embedding space, becomes anomalous:

$$\left\langle r^2(T) \right\rangle \sim T^{2/d_w}, \tag{1.117}$$

with a gap exponent $d_w > 2$. This slowing down of the transport is caused by the fractal ramification of the system. The gap exponent d_w is related to

the spectral dimension d_s and the intrinsic Hausdorff dimension d_h by

$$d_s = \frac{2d_h}{d_w}. \tag{1.118}$$

If the diffusion law is not anomalous, we have $d_w = 2$ and $d_s = d_h$, analogous to $d_s = d$ for diffusion on fixed smooth d-dimensional geometries. For a review of diffusion on fractals see [40].

While the above concepts are in principal valid for Euclidean quantum gravity in any dimension d, let us now specialize to two dimensions. The starting point is again the partition function (1.98) for two-dimensional quantum gravity coupled to D Gaussian matter fields X_μ for fixed volume V of spacetime. We use (1.100) to define $\langle X^2 \rangle_V$. The Gaussian action implies that:

$$\begin{aligned}
\langle X^2 \rangle_V &= \frac{1}{DV} \frac{\partial}{\partial \omega} \left\langle e^{\omega \int_M d^2\xi \sqrt{g}\ X_\mu^2(\xi)} \right\rangle_V \Big|_{\omega=0} \\
&= \frac{1}{DVZ(V)} \frac{\partial}{\partial \omega} \int \mathcal{D}[g_{\mu\nu}]_V \left(\det{}'(-\Delta_g - \omega)\right)^{-D/2} \Big|_{\omega=0} \\
&= \frac{1}{2VZ(V)} \int \mathcal{D}[g_{\mu\nu}]_V \left(\det{}'(-\Delta_g)\right)^{-D/2} \mathrm{Tr}' \left[\frac{1}{-\Delta_g}\right] \\
&= \frac{1}{2V} \left\langle \mathrm{Tr}' \left[\frac{1}{-\Delta_g}\right] \right\rangle_V .
\end{aligned} \tag{1.119}$$

By inserting (1.111) and (1.116) into this formula we get:

$$\begin{aligned}
\langle X^2 \rangle_V &= \frac{1}{2V} \left\langle \int_0^\infty dT \int_M d^2\xi \sqrt{g}\ K'_g(\xi, \xi; T) \right\rangle_V \\
&= \frac{1}{2} \int_0^\infty dT\ RP'_V(T) = \frac{1}{2} \int_0^\infty dT\ \frac{1}{T^{d_s/2}} F\left(\frac{T}{V^{2/d_s}}\right) \\
&\sim V^{\frac{2}{d_s}-1},
\end{aligned} \tag{1.120}$$

for $V \to \infty$. Comparing this with (1.99) we conclude:

$$\frac{1}{d_s} = \frac{1}{D_H} + \frac{1}{2}. \tag{1.121}$$

Using the result $D_H = \infty$ from the preceeding section we arrive at:

$$d_s = 2, \text{ for all } D \leq 1. \tag{1.122}$$

Strictly speaking, this assumes $d_s \leq 2$. However, at $D = -\infty$ a fixed geometry implies $d_s = 2$ and $D_H = \infty$, in agreement with (1.121), and we expect that a saddle point calculation around $D = -\infty$ is reliable, compare

[36]. Therefore d_s equals 2 in a neighbourhood of $D = -\infty$. The average over all geometries in (1.114) includes many degenerate geometries, for which $D_H < \infty$, for example the branched polymer-like geometries which are discussed in section 2.1.4. Thus, a priori we would expect that under the average over fluctuating geometries the spectral dimension decreases. Therefore it is reasonable to assume that $d_s \leq 2$. [2]

Thus we have shown, that the spectral dimension of two-dimensional quantum gravity coupled to D Gaussian fields equals two for $D \leq 1$ [1]. Our derivation relies on the scaling form (1.116) of the averaged return probability $RP'_V(T)$ and on properties of Gaussian matter fields. To corroborate the result (1.122), let us present another derivation completely within Liouville theory[3]. Let us define an observable

$$\mathcal{O}(g) = \int_M d^2\xi_0 \sqrt{g}\, \frac{1}{-\Delta_g} \frac{1}{\sqrt{g(\xi)}} \delta(\xi_0 - \xi)\Big|_{\xi=\xi_0}. \tag{1.123}$$

Under a rescaling of the metric, $\mathcal{O}$ behaves like $\mathcal{O}(\lambda g) = \lambda \mathcal{O}(g)$, that means it is a correlator with conformal weight $(-1,-1)$. Therefore we deduce from (1.96) that

$$\langle \mathcal{O}(g) \rangle_{\lambda V} = \lambda \langle \mathcal{O}(g) \rangle_V, \tag{1.124}$$

or equivalently:

$$\left\langle \frac{1}{V} \mathcal{O}(g) \right\rangle_{\lambda V} = \langle \mathcal{O}(g) \rangle_V. \tag{1.125}$$

On the other hand we can write:

$$\begin{aligned}
\left\langle \frac{1}{V} \mathcal{O}(g) \right\rangle_V &= \left\langle \frac{1}{V} \int_M d^2\xi_0 \sqrt{g(\xi_0)}\, \frac{1}{-\Delta_g} \frac{1}{\sqrt{g(\xi)}} \delta(\xi_0 - \xi)\Big|_{\xi=\xi_0} \right\rangle_V \\
&= \left\langle \frac{1}{V} \int_M d^2\xi_0 \sqrt{g(\xi_0)}\, \int_0^\infty dT\, e^{T\Delta_g} \frac{1}{\sqrt{g(\xi)}} \delta(\xi_0 - \xi)\Big|_{\xi=\xi_0} \right\rangle_V \\
&= \int_0^\infty dT\, RP'_V(T) \sim V^{\frac{2}{d_s}-1},
\end{aligned} \tag{1.126}$$

using the scaling assumption (1.116). From (1.125) and (1.126) it follows that $d_s = 2$ in Liouville theory for all types of matter coupled to gravity

[2] If $d_s > 2$, the integral in (1.120) diverges at 0, and we have to introduce a cut-off ϵ for small times T. Then it is convenient to consider $\langle (X^2)^n \rangle_V \sim V^{2n/D_H}$ instead, with $n = [d_s/2] + 1$ for non-integer d_s. Then the leading large V behaviour will be $V^{2n/d_s - 1}$, and we get $\frac{1}{d_s} = \frac{1}{D_H} + \frac{1}{2n}$.

[3] Thanks to Jakob L. Nielsen for showing me this argument.

with central charge $D \leq 1$, at which point the continuum calculations break down.

This situation is very remarkable: A generic geometry in the functional integral (1.98) is a typical fractal, when looked at in the usual way by computing the fractal dimension d_h from measurements of the circumference $\langle l(g,R)\rangle_V$ of circles versus the geodesic distance R. Also the gap exponent d_w is anomalous and larger than two. It is exactly equal to the intrinsic Hausdorff dimension and consequently, the spectral dimension d_s equals two. It takes the same value as for smooth two-dimensional geometries.

The spectral dimension has received considerable attention also in numerical studies of the fractal structure of two-dimensional quantum gravity. In [39] the return probability is computed for pure gravity and for gravity coupled to matter with central charge $\frac{1}{2}$ and 1. They find that their results are consistent with $d_s = 2$. In [38] d_s is measured for central charge -2, 0, $\frac{1}{2}$ and $\frac{4}{5}$. They get the results $d_s = 2.00(3)$, $d_s = 1.991(6)$, $d_s = 1.989(5)$ and $d_s = 1.991(5)$, respectively[4].

For $D > 1$ it is generally believed that two-dimensional quantum gravity is in a branched polymer phase, see [17, 18] for recent analytical and [41] for some numerical evidence. The Gaussian fields define an embedding of these polymers in $\mathbb{R}^D$. The extrinsic Hausdorff dimension of generic branched polymers is $D_H = 4$, thus we conclude from (1.121) that the spectral dimension equals $\frac{4}{3}$, the famous Alexander-Orbach value [42]. The value $d_s = \frac{4}{3}$ and formula (1.121) for branched polymers have been derived in [43] by different methods, see also [44] for a recent complete proof of $d_s = \frac{4}{3}$. Furthermore, for the multicritical branched polymers [21] it is known that $D_H = \frac{2m}{m-1}, m = 2, 3, \ldots$, where $m = 2$ for the ordinary branched polymers. Thus we obtain $d_s = \frac{2m}{2m-1}$, in agreement with the analysis in [45]. For $m \to \infty$ the multicritical branched polymers approach ordinary random walks, for which $D_H = 2$ and $d_s = 1$.

Let us in the end comment on a subtlety in the definition (1.114) of the averaged return probability $RP'_V(T)$. We have defined it as the quantum average of the return probability for fixed geometry over fluctuating geometries. Instead, we could have defined it as the limit for $R \to 0$ of

$$K'_V(R,T) = \frac{1}{G(V,R)} \int \mathcal{D}[g_{\mu\nu}]_V \, e^{-S_{\text{eff}}(g)} \times \int_M d^2\xi_1 \sqrt{g} \int_M d^2\xi_2 \sqrt{g} \, \delta(R - d_g(\xi_1,\xi_2)) K'_g(\xi_1,\xi_2;t), \tag{1.127}$$

[4]Thanks to Konstantinos N. Anagnostopoulos for showing me these data prior to the publication of [38]

where $G(V,R) = V\langle l(g,R)\rangle_V Z(V)$ is the partition function for universes with two marked points separated by a geodesic distance R ("two-point function") as defined (up to matter fields) in (1.25), see also (1.35). $K'_V(R,T)$ is the average probability distribution for diffusing a geodesic distance R in the time T. It is natural to identify $RP'_V(T)$ with $K'_V(0,T)$. However it is unknown whether the limit $R \to 0$ of (1.127) commutes with the functional integration over geometries. Actually, for the two-point function $G(V,R)$ it is easy to see that this is not the case. If the limits do not commute there are two inequivalent definitions of the return probability $RP'_V(T)$. However, the scaling form of $K'_V(0,T)$ is the same as the scaling form (1.116) of $RP'_V(T)$ [46]. Indeed, we have

$$\int_0^\infty dR\, \langle l(g,R)\rangle_V \; K'_V(R,T) = 1, \tag{1.128}$$

and using $\dim[R] = \dim[V^\nu]$ we arrive at the scaling form

$$K'_V(R,T) = \frac{1}{V} P\left(\frac{R}{V^\nu}, \frac{T}{V^\lambda}\right), \tag{1.129}$$

where λ is chosen such that $\frac{T}{V^\lambda}$ is dimensionless. Expanding the return probability $K'_V(0,T)$ around $T = 0$ we get

$$K'_V(0,T) \sim \frac{V^{\frac{\lambda d_s}{2}-1}}{T^{d_s/2}}, \text{ for } T \sim 0, \tag{1.130}$$

by definition of d_s. If this short time asymptotic stays nonzero and finite for infinite volume V we get $\lambda = \frac{2}{d_s}$, and $K'_V(0,T)$ scales as

$$K'_V(0,T) = \frac{1}{T^{d_s/2}} \tilde{F}\left(\frac{T}{V^{2/d_s}}\right), \tag{1.131}$$

which has the same form as the scaling behaviour of $RP'_V(T)$.

1.3.3 Intrinsic Hausdorff dimension

The intrinsic Hausdorff dimension d_h of two-dimensional quantum gravity has been defined in section 1.1.3. One way is to measure the average circumference $\langle l(g,R)\rangle_V$ of circles with a small radius R, which scales as $\langle l(g,R)\rangle_V \sim R^{d_h - 1}$. This is analogous to the classical "clock and rod" procedure to study the geometrical properties of spacetime. Alternatively one can define the fractal dimension d_H as the scaling exponent of the average volume of the universes: $\langle V\rangle_R \sim R^{d_H}$, compare (1.32). In two-dimensional

quantum gravity the first "local" and the second "global" concept of the intrinsic Hausdorff dimension agree. For pure gravity, we have $d_H = 4$, see (1.41).

The intrinsic Hausdorff dimension has been computed in [33] by studying the diffusion equation in Liouville theory. Although the mathematical status of this derivation is unclear and although it is not consistent with the derivation of the spectral dimension in section 1.3.2, the result is valid for a number of special cases and agrees with many numerical simulations.

In [33] the mean square distance $\left\langle R^2(T)\right\rangle_V$ of diffusion in the fluctuating space time is defined as:

$$\begin{aligned}\left\langle R^2(T)\right\rangle_V \;\equiv\; & \frac{1}{VZ(V)}\int \mathcal{D}[g_{\mu\nu}]_V\; e^{-S_{\text{eff}}(g)}\int_M d^2\xi_1\sqrt{g}\int_M d^2\xi_2\sqrt{g} \\ & \times d_g^2(\xi_1,\xi_2)K'_g(\xi_1,\xi_2;T). \end{aligned} \tag{1.132}$$

$\left\langle R^2(T)\right\rangle_V$ scales with the volume as

$$\left\langle R^2(T)\right\rangle_V \sim V^{\frac{2}{d_h}}, \tag{1.133}$$

which yields yet another way to define the intrinsic Hausdorff dimension d_h. On the other hand, the mean square distance for diffusion on a fixed smooth geometry $[g_{\mu\nu}]$ with the initial condition (1.109) can be expanded for small times T as [47]

$$\int_M d^2\xi\sqrt{g}\; d_g^2(\xi_0,\xi)K'_g(\xi_0,\xi;T) = 4T - \frac{2}{3}T^2\mathcal{R}(\xi_0) + O(T^3), \tag{1.134}$$

with the curvature scalar $\mathcal{R}$. If we assume that this expansion commutes with the functional average over geometries in (1.132), we get

$$\left\langle R^2(T)\right\rangle_V \sim \text{const}\;\cdot T, \text{ for small times } T. \tag{1.135}$$

The dimension of T can be computed from Liouville theory by expanding the (unnormalized) return probability for small times to observe how this scales under the transformation $V \to \lambda V$:

$$\begin{aligned}\left\langle \int_M d^2\xi_0\sqrt{g}\; K'_g(\xi_0,\xi_0;T)\right\rangle_{\lambda V} &= \left\langle \int_M d^2\xi_0\sqrt{g}\; e^{T\Delta_g}\Psi(\xi,0)\Big|_{\xi=\xi_0}\right\rangle_{\lambda V} \\ &= \left\langle \int_M d^2\xi_0\sqrt{g}\;\left[\Psi(\xi_0,0) + T\Delta_g\Psi(\xi,0)\Big|_{\xi=\xi_0} + O(T^2)\right]\right\rangle_{\lambda V} \\ &= 1 + T\left\langle \int_M d^2\xi_0\sqrt{g}\;\Delta_g\frac{1}{\sqrt{g(\xi)}}\delta(\xi-\xi_0)\Big|_{\xi=\xi_0}\right\rangle_{\lambda V} + O(T^2) \\ &\overset{(1.96)}{=} 1 + T\,\lambda^{\frac{\beta}{\alpha}} + O(T^2), \end{aligned} \tag{1.136}$$

where β is computed for $\Delta_0 = 2$. This calculation is based on the assumption that the expansion of the exponential commutes with the functional integration over geometries. Then we have:

$$\dim[T] = \dim[V^{2/d_H}] = \dim[V^{-\beta/\alpha}], \tag{1.137}$$

and thus

$$d_H = -\frac{2\alpha}{\beta} = 2\,\frac{\sqrt{25-D}+\sqrt{49-D}}{\sqrt{25-D}+\sqrt{1-D}}, \tag{1.138}$$

for matter with central charge D coupled to two-dimensional gravity. For $D = 0$, that means in the case of pure gravity, (1.138) predicts $d_H = 4$, in agreement with (1.41). In the semiclassical limit $D \to -\infty$ we expect $d_H = 2$, which is also correctly predicted by (1.138).

The validity of (1.138) has been widely discussed. The main assumption of the derivation sketched above is that one is allowed to interchange the expansion of $e^{T\Delta_g}$ with the functional integration in (1.136). Further input is the scaling assumption for $\langle R^2(T)\rangle_V$. In principle the same result for the intrinsic Hausdorff dimension should be obtained if one uses some power of the Laplace operator. However this is not true and results obtained from higher orders of the expansion of $e^{T\Delta_g}$ contradict the result from the first order.

An alternative theoretical prediction based on the transfer matrix method has been given in [34, 48, 49]. They get:

$$d_H = -\frac{2}{\gamma} = \frac{24}{1-D+\sqrt{(25-D)(1-D)}}. \tag{1.139}$$

For $D = 0$ this yields again $d_H = 4$, while for $D \to -\infty$ (1.139) predicts $d_H = 0$.

To settle this question, the authors of [50, 51] performed a high precision Monte Carlo analysis for $D = -2$. Their result is $d_H = 3.574(8)$, in agreement with the value $d_H = 3.562$ from (1.138) and in clear disagreement to $d_H = 2$ from (1.139). This clearly rules out the validity of (1.139) for matter with $D < 0$ coupled to gravity.

However, for unitary matter with $D > 0$ coupled to gravity the situation is less clear. Most simulations report values $d_H \approx 4$ [52, 39, 53], and a back-reaction of the matter on the structure of the quantum spacetime cannot be observed. However, values extracted from the scaling of the matter correlation functions are higher, though less reliable. All data are consistent with the conjecture [52, 39]:

$$d_H = 4, \text{ for } 0 \le D \le 1, \tag{1.140}$$

$D=-5$	$D=-2$	$D=0$	$D=\frac{1}{2}$	$D=\frac{4}{5}$	$D=1$	ref.
3.236	3.562	4	4.212	4.421	4.828	(1.138)
1.236	2	4	6	10	∞	(1.139)
3.36(4)	3.574(8)	4.05(15)	4.11(10)	4.01(9)	3.8–4.0	[54, 55]
			3.96–4.38	3.97–4.39		[55]

Table 1.1: Current status of numerical values for the intrinsic Hausdorff dimension d_H compared to the two different theoretical predictions (1.138) and (1.139). The values in the first part of the table are the theoretical predictions. In the second part of the table we summarize results from the scaling of the two-point function for the random surfaces, while in the last part the Hausdorff dimension is determined from the scaling of the matter correlation functions. See the references for technical details about the simulations and the data analysis.

although the validity of (1.138) for $D > 0$ is not completely ruled out. The current status of the numerical simulations is summarized in table 1.1.

Chapter 2

Discretization of 2d quantum gravity

The formulation of quantum field theory in terms of renormalized Euclidean functional integrals leads to an identification of quantum field theory with statistical mechanics. A key ingredient for this identification is the discretization of spacetime. Then the powerful techniques from the theory of critical phenomena get available and have proved to be invaluable for the understanding of renormalization and of non-perturbative phenomena.

Therefore it is natural to attempt a discretization of the continuum theory of quantum gravity. However, spacetime no longer plays the role of a mere background but is the dynamical variable itself. This, combined with the diffeomorphism invariance of the continuum theory poses a problem for the discretization which has successfully been solved by the method of dynamical triangulations [56, 57, 58]. The main idea is to discretize geometry directly, with no reference to coordinate parametrizations. That means that the functional integral over equivalence classes of metrics on a manifold M is replaced by a finite sum over piecewise linear spaces which are constructed by successively gluing d-simplices together. The fixed edge length of the simplices introduces a reparametrization invariant cutoff into the theory. In this way quantum gravity can be formulated as an ensemble of discrete random manifolds and the machinery of critical phenomena can be applied. The continuum theory is recovered at a critical point of this ensemble. Masses and continuum coupling constants are defined by the approach to the critical point in the scaling limit.

Research in the formulation of classical gravity in purely geometrical terms was initiated in 1961 by T. Regge [59], while the first attempt at a discretization of quantum gravity in the sense of this chapter has been

done by Weingarten in 1980 [60, 61]. A good review with some comments about the historical development of the subject is [62]. Other reviews can be found in [63, 64, 65]. Much of the material presented in this chapter is taken from these articles. In [66] the relation between the matrix-model technique and dynamical triangulation is reviewed.

In the first section of this chapter we introduce the method of dynamical triangulations and discuss the scaling properties of the theory. Branched polymers, which have been mentioned before, are discussed in some detail. In section 2.2 we give a brief review of the matrix-model techniques which have been applied to solve two-dimensional quantum gravity. For more details and further issues we refer to the review articles given above. The explicit solution of discretized two-dimensional quantum gravity shows, that the continuum theory is the scaling limit of dynamical triangulation. Therefore both theories can be identified.

In section 2.3 we outline how the fractal structure and the scaling properties of pure two-dimensional quantum gravity can be obtained.

An alternative method for the discretization of quantum gravity, known as quantum Regge calculus, has been suggested. This is discussed in chapter 3 of this thesis.

2.1 Dynamical triangulation

Dynamical triangulation is a discretization of quantum geometries. No reference to parametrizations has to be made. At the same time it provides a regularization of the continuum theory by introducing an explicit cutoff. In this section we can only outline the beauty and power of dynamical triangulations in two dimensions. We begin with a short introduction into discretization of continuum manifolds and geometries. Then we define the partition function of two-dimensional dynamical triangulation and discuss its scaling properties. The susceptibility exponent γ is characterized through the fractal structure of the random spacetimes. We end this section with a discussion of the scaling properties of the two-point function which are illustrated by means of the model of branched polymers.

2.1.1 Discretization of geometry

Let us define an oriented n-simplex as an $(n+1)$-tuple of points modulo even permutations. Thus a 0-simplex is a point, a 1-simplex is a pair of

points which can be identified with a line, a 2-simplex is a triangle, a 3-simplex a tetrahedron etc. A subsimplex σ' of a simplex σ is defined as a subset of σ: $\sigma' \subseteq \sigma$. By gluing simplices together one gets a simplicial complex. Simplicial complexes have no fixed dimension, that means they are not manifolds. However, their structure is sufficient to define exterior calculus on them, which physically means that scalar fields, gauge fields, antisymmetric tensor fields etc can be defined on them.

A d-dimensional simplicial manifold is a simplicial complex such that the neighbourhood of each point in the complex is homeomorphic to a d-dimensional ball. In one and two dimensions each simplicial manifold can be obtained by gluing pairs of d-simplices along some of their $(d-1)$-edges. In higher dimensions this successive gluing yields in some cases only pseudo-manifolds, in which a neighbourhood of a point can also be homeomorphic to a topologically more complicated object.

The structure of simplicial manifolds corresponds to a discretization of the continuous manifold structure. To define geometrical concepts we have to introduce a metric. A d-dimensional simplicial manifold can be equipped canonically with a Riemannian metric by demanding that the metric is flat inside each d-simplex, continuous when the $(d-1)$-subsimplices are crossed and that each $(d-1)$-subsimplex is a linear flat subspace of this simplex. A simplicial manifold equipped with such a metric is called piecewise linear space.

The canonical metric can be defined by assigning lengths to the N_l links (1-subsimplices) of the simplicial manifold. First note that any d-dimensional simplicial manifold M can be covered with charts (U, ϕ) such that each d-simplex is parametrized by barycentric coordinates. Here

$$U = \{\xi \in \mathbb{R}^d_+ | \xi_1 + \cdots + \xi_d < 1\}, \tag{2.1}$$

and $\phi : U \to M$ is given by

$$\phi(\xi) = \xi_1 y_1 + \ldots + \xi_d y_d + (1 - \xi_1 - \ldots - \xi_d) y_{d+1}, \tag{2.2}$$

where $y_1, \ldots, y_{d+1}$ are the coordinates of the vertices of the simplicial complex which live in some ambient space $\mathbb{R}^n$. Then on each chart the canonical metric is defined by

$$g_{\mu\nu}(\xi) = \frac{\partial \phi}{\partial \xi^\mu} \cdot \frac{\partial \phi}{\partial \xi^\nu}. \tag{2.3}$$

This metric is Euclidean inside each d-simplex and continuous if a $(d-1)$-face is crossed. It is compatible with the manifold structure and can be

expressed solely in terms of the link lengths $l_1, \ldots, l_{N_l}$. For a triangle (2.3) can be written as

$$g_{\mu\nu} = \begin{pmatrix} l_1^2 & \frac{1}{2}(l_1^2 + l_2^2 - l_3^2) \\ \frac{1}{2}(l_1^2 + l_2^2 - l_3^2) & l_2^2 \end{pmatrix}. \tag{2.4}$$

The intrinsic curvature of a two-dimensional piecewise linear space is located at the vertices, that means its 0-subsimplices. The curvature can clearly not reside in the triangles, since they are flat. Furthermore the metric is continuous when crossing the edges of the triangles. Since we define curvature as an intrinsic concept of the piecewise linear space it should be independent of the embedding. We can bend the surface around an edge without changing the geometry – an observation which the German mathematician C. F. Gauss made in his famous paper *Disquisitiones generales circa superficies curva* (1828) and which he called *theorema egregrium*. Therefore it is intuitive that the curvature cannot reside in the edges. Rather, it is concentrated at the vertices or in general at the $(d-2)$-hinges of the piecewise linear space. To each vertex v in the surface we can assign a deficit angle δ_v as the difference between 2π and the sum of the angles meeting at v:

$$\delta_v = 2\pi - \sum_{t: v \in t} \alpha_{v,t}, \tag{2.5}$$

where the sum goes over all triangles t containing v. $\alpha_{v,t}$ is the angle at v inside t. The scalar curvature $\mathcal{R}_v$ at v is defined as

$$\mathcal{R}_v = 2\frac{\delta_v}{dA_v}, \; dA_v = \frac{1}{3}\sum_{t:\, v \in t} A_t, \tag{2.6}$$

where A_t is the area of the triangle t. The scalar curvature $\mathcal{R}$ in two dimensions equals two times the Gaussian curvature. With these definitions we finally have:

$$\text{total area of the surface} = \sum_{v=1}^{N_v} dA_v, \tag{2.7}$$

and the Gauss-Bonnet theorem is now valid in the form

$$\sum_{v=1}^{N_v} \mathcal{R}_v dA_v = 4\pi\chi, \tag{2.8}$$

which is equivalent to the polyeder formula $N_t - N_l + N_v = \chi$ by Descartes and Euler.

2.1.2 Definition of the model

The original proposal of Regge [59] consists in constructing a sequence of piecewise linear spaces to approximate a given smooth manifold M such that for any sufficiently smooth function f

$$\sum_{\nu} dA_\nu f_\nu \to \int_M dA(\xi)\ f(\xi), \tag{2.9}$$

where f_ν is the value of the function f at the vertex ν. The applicability of this point of view in the quantum case will be discussed in chapter 3. Here we take a somewhat different point of view. Our aim is to integrate over all equivalence classes of metrics on a given manifold M. We introduce a reparametrization invariant cutoff by assigning the length a to each edge of piecewise linear spaces. We construct all possible piecewise linear spaces by gluing equilateral triangles together. In this case the formulas above simplify considerably. Let n_ν be the order of the vertex ν. Then we have:

$$dA_\nu = \frac{\sqrt{3}}{12} a^2 n_\nu, \quad \mathcal{R}_\nu dA_\nu = \frac{2\pi}{3}(6 - n_\nu). \tag{2.10}$$

For simplicity we rescale a^2 by $\frac{\sqrt{3}}{4}$ and then set $a = 1$. (2.10) shows that two triangulations which cannot be mapped onto each other by a relabelling of the vertices lead to different local curvature assignments. Thus they define different metric structures. The set of combinatorially non-equivalent triangulations defines a grid in the space of diffeomorphism classes of metrics. The proposal described in this chapter relies on the hope that this grid gets uniformly dense in the limit $a \to 0$.

To define a regularized theory of quantum gravity we replace the action $S(g, \Lambda) = \Lambda V_g$ in (1.6) by

$$S_T(\mu) = \mu N_t, \tag{2.11}$$

where μ is the bare cosmological constant. The integration over equivalence classes of metrics on M is replaced by a summation over all non-isomorphic equilateral triangulations T with the topology of M. In the case of matter coupled to quantum gravity, the matter fields X can be defined on the vertices, the links or on the triangles. In general the action S_{matter} and the functional integration over the matter fields will depend on the triangulation T. The discretized version of the partition function can thus be written as

$$Z(\mu) = \sum_{T \in \mathcal{T}} \frac{1}{C_T} e^{-S_T(\mu)} \int \mathcal{D}_T[X]\ e^{-S_{\text{matter}}(T,X)}, \tag{2.12}$$

where C_T denotes the symmetry factor of the triangulation T which is equal to the order of the automorphism group of T. The summation goes over a suitable class $\mathcal{T}$ of abstract triangulations T defined by their vertices and a connectivity matrix. Appealing to universality details of the chosen class of triangulations such as whether closed two-loops are allowed or not should not be important for the theory. This has been verified a posteriori.

For simplicity we will ignore any possible matter fields coupled to gravity in the following equations.

Then we can write:

$$Z(\mu) = \sum_{N=1}^{\infty} e^{-\mu N} Z(N), \quad Z(N) = \sum_{T \in \mathcal{T}_N} \frac{1}{C_T}, \tag{2.13}$$

where $\mathcal{T}_N$ denotes the subset of triangulations with N triangles of $\mathcal{T}$. Expectation values of observables $\mathcal{O}$ are defined as:

$$\langle \mathcal{O} \rangle_\mu = \frac{1}{Z(\mu)} \sum_{T \in \mathcal{T}} \frac{1}{C_T} e^{-S_T(\mu)} \mathcal{O}(T), \tag{2.14}$$

$$\langle \mathcal{O} \rangle_N = \frac{1}{Z(N)} \sum_{T \in \mathcal{T}_N} \frac{1}{C_T} \mathcal{O}(T), \tag{2.15}$$

in the "grand-canonical" ensemble with fixed cosmological constant and in the "canonical" ensemble with fixed volume respectively. μ can be understood as the chemical potential for adding new triangles to the surfaces.

$Z(N)$ can be interpreted as the number of triangulations in $\mathcal{T}_N$. This number is exponentially bounded:

$$Z(N) = e^{\mu_c N} N^{\gamma - 3} \left(1 + O(1/N)\right). \tag{2.16}$$

The proof for spherical topology has been given in [67]. For a general proof by purely combinatorial methods in the spirit of Tutte see [62]. It can be proved that the critical point μ_c does not depend on the topology.

That means that the statistical ensemble defined by (2.12) has a critical point μ_c. $Z(\mu)$ is analytical for $\mu > \mu_c$ and contains non-analytical parts at the critical point. The latter have our biggest interest since they are the universal parts given by the large N behaviour of $Z(N)$. The continuum limit should be defined as $\mu \to \mu_c$. Close to the critical point we have:

$$Z(\mu) = (\mu - \mu_c)^{2-\gamma} + \text{less singular terms}, \tag{2.17}$$

from a discrete Laplace transformation of (2.16). Many analytical and numerical investigations have revealed that γ assumes the same values as in the

continuum theory of two-dimensional quantum gravity and is given by (1.14). Thus the susceptibility exponent γ can also be characterized as the subleading power correction of the exponentially growing number of triangulations for a fixed topology. Furthermore, γ can be characterized by the branching ratio into minimal bottleneck baby universes [68], that means parts of the universe which are connected to the rest by a three-loop of links which do not form a triangle, see figure 2.1. The smaller part is then called minimal



Figure 2.1: Branching of a minimal bottleneck baby universe from the larger parent.

bottleneck baby universe and the larger part is called parent. In the ensemble of universes with volume N the average number $\langle \mathcal{N}(V) \rangle_N$ of baby universes with volume V is given by

$$\langle \mathcal{N}(V) \rangle_N \sim \frac{3!}{Z(N)} V Z(V)\ (N-V) Z(N-V). \tag{2.18}$$

The factors $Z(V)$ and $Z(N-V)$ are the weights of the baby universe and of the parent respectively. The volume factors V and $N-V$ reflect the fact that the baby universe can be attached at any triangle. Finally, $3!$ is the number of ways the two boundaries of the parent and of the minimal bottleneck baby universe can be glued together. (2.18) is valid in the generic case where there are no additional symmetries. Assuming that $Z(N)$ is given by (2.16) we get:

$$\langle \mathcal{N}(V) \rangle_N \sim N V^{\gamma-2} \left(1 - \frac{V}{N}\right)^{\gamma-2}. \tag{2.19}$$

For $V \ll N$ this reduces to $\langle \mathcal{N}(V) \rangle_N \sim N V^{\gamma-2}$. It is interesting to note that γ is a function of the matter coupled to gravity. Thus it describes an

aspect of the fractal structure of quantum spacetime that originates from the back-reaction of the matter on gravity.

2.1.3 The two-point function

In analogy to the continuum formalism in chapter 1 we can define a two-point function which in a natural way contains information about the fractal structure of the random surfaces.

In principle, we could define a geodesic distance by using the canonical metric described in section 2.1.1. Alternatively, the use of simplified definitions has been suggested. The geodesic distance between two links l_1 and l_2 is defined as the length of the shortest path of triangles connecting l_1 and l_2. The geodesic distance between a link l_1 and a set of links $\mathcal{L}$ is defined as the minimum of the geodesic distances between l_1 and the elements of $\mathcal{L}$. Furthermore we define the geodesic distance between a loop $\mathcal{L}_1$ and a loop $\mathcal{L}_2$ to be r if all links of $\mathcal{L}_1$ lie a geodesic distance r from the loop $\mathcal{L}_2$. Note that this definition is not symmetric in $\mathcal{L}_1$ and $\mathcal{L}_2$. Similarly we could have defined the geodesic distance as the distance between vertices or between triangles.

Then the two-point function is defined as:

$$G(\mu, r) = \sum_{T \in \mathcal{T}(2,r)} e^{-\mu N_t} = \sum_{N=1}^{\infty} e^{-\mu N} G(N, r), \tag{2.20}$$

$$G(N, r) = \sum_{T \in \mathcal{T}_N(2,r)} 1, \tag{2.21}$$

where $\mathcal{T}(2, r)$ denotes the class of triangulations T with two marked links separated a geodesic distance r, and $\mathcal{T}_N(2, r)$ the subclass where all triangulations consist of N triangles. Note that r is an integer in units of the lattice spacing. $G(\mu, r)$ is the discretized form of $G(\Lambda, R)$ in (1.23), while $G(N, r)$ is the discretized form of $G(V, R)$ in (1.25).

An important property of the two-point function is:

$$G(\mu, r_1 + r_2) \geq \text{const} \times G(\mu, r_1)\ G(\mu, r_1), \tag{2.22}$$

which follows from the fact that each term on the right hand side uniquely defines a term on the left hand side. This is demonstrated in figure 2.2. From this one can conclude that the limit

$$\lim_{r \to \infty} \frac{-\log G(\mu, r)}{r} = m(\mu) \tag{2.23}$$



Figure 2.2: Two surfaces with two marked links separated a geodesic distance r_1 and r_2 respectively can be glued together after cutting open one of the marked links each to form a surface with two marked links separated a geodesic distance $r_1 + r_2$.

exists with $m(\mu) \geq 0$ and $m'(\mu) > 0$, for $\mu > \mu_c$. Thus the two-point function falls off exponentially for $r \to \infty$. These relations are by now well known. However, the complete proofs require some technical arguments, compare [62].

Below we will show that a continuum limit of the discretized theory can only exist if $m(\mu)$ scales to zero as $\mu \to \mu_c$. Let us assume that this is the case:

$$m(\mu) \sim (\mu - \mu_c)^{\nu}, \text{ for } \mu \to \mu_c. \tag{2.24}$$

In general we expect that close to the critical point the exponential decay of $G(\mu, r)$ turns over into a power law. More precisely:

$$G(\mu, r) \sim e^{-m(\mu)r}, \text{ for } 1 \ll m(\mu)r, \tag{2.25}$$

$$G(\mu, r) \sim r^{1-\eta}, \text{ for } m(\mu) \ll m(\mu)r \ll 1. \tag{2.26}$$

The exponent η is called the anomalous dimension in analogy to ordinary statistical systems. Furthermore we can define the susceptibility $\chi(\mu)$ in the discretized ensemble as

$$\chi(\mu) = \sum_{r=1}^{\infty} G(\mu, r), \tag{2.27}$$

Close to the critical point, where triangulations with large N_t dominate and symmetry factors play no role, we have

$$\chi(\mu) \sim \frac{\partial^2 Z(\mu)}{\partial \mu^2} \sim (\mu - \mu_c)^{-\gamma}. \tag{2.28}$$

Similar to chapter 1 we can introduce several concepts of fractal dimensions to characterize the geometrical structure of the two-dimensional quantum spacetime. The Hausdorff dimension d_H in the grand-canonical ensemble is defined by

$$\langle N \rangle_r \sim r^{d_H}, \text{ for } r \to \infty \text{ and } m(\mu) r = \text{const}, \tag{2.29}$$

where the average $\langle N \rangle_r$ is given by

$$\langle N \rangle_r = \frac{1}{G(\mu, r)} \sum_{T \in \mathcal{T}(2,r)} N_t e^{-\mu N_t} = -\frac{\partial \log G(\mu, r)}{\partial \mu}. \tag{2.30}$$

If we perform the derivative under the constraint in (2.29) it follows that

$$\langle N \rangle_r \sim m'(\mu) r \sim r^{\frac{1}{\nu}}. \tag{2.31}$$

Thus the Hausdorff dimension d_H is related to the scaling exponent ν by:

$$\nu = \frac{1}{d_H}. \tag{2.32}$$

With (2.28) we can derive Fisher's scaling relation

$$\gamma = \nu(2 - \eta), \tag{2.33}$$

which relates the critical exponents defined above. From the long distance behaviour of $G(\mu, r)$ we can compute the long distance behaviour of $G(N, r)$ by a saddle point calculation:

$$G(N, r) \sim e^{-c\left(\frac{r}{N^\nu}\right)^{\frac{1}{1-\nu}}}, \tag{2.34}$$

with $c = \left(\frac{1}{\nu} - 1\right) \nu^{\frac{1}{1-\nu}}$.

Analogously to the continuum formalism in chapter 1 we can define a Hausdorff dimension d_h by the short distance scaling of the two-point function in the canonical ensemble with fixed number N of triangles. For $r = 0$, $G(N, 0)$ is the one-point function and behaves as

$$G(N, 0) \sim e^{\mu_c N} N^{\gamma - 2}. \tag{2.35}$$

This is because for large N the one-point function is proportional to $NZ(N)$ since it counts triangulations with one marked link.

Now let $r \ll N^{1/d_h}$ and count the number $n(r)$ of triangles which lie a geodesic distance r away from a marked link. The average of $n(r)$ in the

ensemble of surfaces with one marked link defines the Hausdorff dimension d_h by

$$\langle n(r)\rangle_N \sim \frac{G(N,r)}{G(N,0)} \sim r^{d_h-1}, \text{ for } 1 \ll r \ll N^{1/d_h}. \tag{2.36}$$

The first relation follows from the definition of $\langle n(r)\rangle_N$. Together with (2.35) we conclude:

$$G(N,r) \sim r^{d_h-1} N^{\gamma-2} e^{\mu_c N}, \text{ for } 1 \ll r \ll N^{1/d_h}. \tag{2.37}$$

Finally, the short distance behaviour of $G(\mu,r)$ follows by a discrete Laplace transformation. Close to the critical point we get:

$$G(\mu,r) \sim r^{\gamma d_h-1}. \tag{2.38}$$

Together with (2.26) this proves again Fisher's scaling relation. Furthermore it turns out that both definitions of the intrinsic Hausdorff dimension are equivalent in two-dimensional quantum gravity. However, counter examples exist, as will be demonstrated in the next section.

The mass $m(\mu)$ determines the scaling of the discretized theory. To see how continuum expressions can be approached in the discretized theory let us reintroduce dimensions. The renormalized cosmological constant Λ in terms of the bare cosmological constant μ is defined by:

$$\mu - \mu_c = \Lambda a^2, \text{ that means } a(\mu) \sim (\mu-\mu_c)^{1/2}. \tag{2.39}$$

If the mass shall survive in the continuum limit, it has to be introduced as:

$$M = m(\mu)\ a(\mu)^{-2\nu} = c\Lambda^{\nu}, \tag{2.40}$$

with a constant c. If the two-point function shall survive, we must have

$$m(\mu)r = MR, \tag{2.41}$$

where the continuum parameter M and R are held fixed and the number of steps r goes to infinity, that means:

$$R = r\ a(\mu)^{2\nu}. \tag{2.42}$$

From (2.25) and (2.26) we see that the continuum two-point function is given by:

$$G(\Lambda,R) = \lim_{\mu\to\mu_c} a(\mu)^{2\nu(1-\eta)} G(\mu,r), \quad m(\mu)r = MR. \tag{2.43}$$

This scaling form can be explicitly verified by an exact calculation of $G(\mu,r)$ for pure gravity [22, 23], compare section 2.3.

2.1.4 Branched polymers

An interesting simple example for the scenario described above is provided by the model of branched polymers which is important in the understanding of the phases of two-dimensional quantum gravity. The critical indices have been computed in detail in [69, 21]. The partition function is defined as

$$Z(\mu) = \sum_{BP} \frac{1}{C_{BP}} e^{-\mu N_l} \prod_{v \in BP} f(n_v), \tag{2.44}$$

where the sum goes over all branched polymers, that means all connected planar tree graphs. N_l denotes the number of links in a branched polymer. The product goes over all vertices in the tree graphs and n_v is the number of links joining at the vertex v, called the order of v. Usually, the (unnormalized) branching weight $f(n)$ is a non-negative function of the order of the vertices. Finally the symmetry factor C_{BP} is chosen such that rooted branched polymers, that means with the first link marked, are counted only once.

The one-point function $G(\mu)$ is defined similarly as the sum over the rooted branched polymers. In this case, the symmetry factor drops out.



Figure 2.3: Illustration of the self-consistent equation for rooted branched polymers. The first link gives a factor $e^{-\mu}$. At the vertices the branching weight has to be included.

From figure 2.3 we follow that $G(\mu)$ satisfies the equation

$$G(\mu) = e^{-\mu} \left(1 + f(2)G(\mu) + f(3)G(\mu)^2 + \dots\right). \tag{2.45}$$

We can solve this relation for e^{μ} as a function of $G(\mu)$:

$$e^{\mu} = \frac{1 + f(2)G(\mu) + f(3)G(\mu)^2 + \dots}{G(\mu)} \equiv \frac{\mathcal{F}(G)}{G} \equiv F(G). \tag{2.46}$$



Figure 2.4: Graphical solution of (2.46). The critical point is identified as the minimum of e^{μ} as a function of G.

This equation is illustrated in figure 2.4. The critical point is at the minimum of $F(G)$. Since all $f(n)$ are positive the minimum is unique and satisfies $F'(G(\mu_c)) = 0$ and $F''(G(\mu_c)) > 0$. Therefore we have:

$$G(\mu) \sim G(\mu_c) - c(\mu - \mu_c)^{\frac{1}{2}}, \text{ for } \mu \to \mu_c, \tag{2.47}$$

where c is some constant. Since $G(\mu)$ is the one-point function we would have expected a behaviour $G(\mu) \sim (\mu - \mu_c)^{1-\gamma}$. Thus we conclude that the generic value of γ for branched polymers is $\frac{1}{2}$.

If we allow some of the values $f(n)$ to be negative, we can fine tune the $f(n)$ such that the minimum of $F(G)$ satisfies

$$F^{(k)}(G(\mu_c)) = 0, \text{ for } k = 1, \ldots, m-1, \text{ and } F^{(m)}(G(\mu_c)) \neq 0. \tag{2.48}$$

This model is called m'th multicritical branched polymer model. For $m = 2$ we simply recover the ordinary branched polymer model. The critical behaviour of the one-point function is changed to

$$G(\mu) \sim G(\mu_c) - c(\mu - \mu_c)^{\frac{1}{m}}, \text{ for } \mu \to \mu_c, \tag{2.49}$$

and we get $\gamma = \frac{m-1}{m}$.

To define the two-point function $G(\mu, r)$ for branched polymers, let the geodesic distance between two marked vertices x and y be the minimal length of paths between the two points. This is of course unique. The two-point function is defined as the partition function of branched polymers with the constraint that there are two marked points separated a geodesic distance r.

The path between the marked points can be viewed as a random walk where at each vertex a rooted branched polymer can grow out. If we think only in terms of intrinsic geometrical properties this leads to [21]:

$$G(\mu, r) \sim (e^{-\mu}\mathcal{F}'(G(\mu)))^r = e^{-m(\mu)r}, \tag{2.50}$$

from which we conclude, using $e^{-\mu}\mathcal{F}'(G(\mu)) = 1 - \frac{G(\mu)}{\chi(\mu)}$, that

$$m(\mu) = \kappa(\mu - \mu_c)^{\frac{m-1}{m}}, \tag{2.51}$$

in the general case for $m \geq 2$, with a positive constant κ. Therefore it follows with (2.32) that the intrinsic Hausdorff dimension d_H equals $\frac{m}{m-1}$. For $m = 2$ we have $d_H = 2$ and the branched polymers have the same intrinsic dimension as smooth two-dimensional manifolds. For $m \to \infty$, d_H approaches one, in agreement with the fact that in this limit the branched polymers approach ordinary random walks. The two-point function $G(\mu, r)$ is given by

$$G(\mu, r) = \text{const} \times e^{-\kappa r(\mu-\mu_c)^{\frac{m-1}{m}}}. \tag{2.52}$$

We can also compute the canonical intrinsic Hausdorff dimension d_h for branched polymers defined in the ensemble of branched polymers with a fixed number N of links. The volume $\langle n(r)\rangle_N$ of a spherical shell of geodesic radius r is defined in analogy to (1.35) and (2.36). $Z(N)$ for branched polymers is given by an inverse Laplace transformation of (2.44) and scales as $Z(N) \sim e^{\mu_c N} N^{\gamma-3}$. Similarly the two-point function $G(N, r)$ for graphs with fixed volume N is given by an inverse Laplace transformation of $G(\mu, r)$. Inserting the scaling behaviour and taking only the leading orders we find for small r [51]:

$$\langle n(r)\rangle_N \sim \frac{r}{N^{1-\frac{2}{m}}}, \text{ for } N^{1-\frac{2}{m}} \ll r \ll N^{1-\frac{1}{m}}. \tag{2.53}$$

This shows that $d_h = 2$ for all values of m. Thus for the generic branched polymers ($m = 2$) the canonical and the grand-canonical definition of the intrinsic Hausdorff dimension give the same result. However, this is not true for the m'th multicritical branched polymer model with $m > 2$, where $d_H = \frac{m}{m-1}$ and $d_h = 2$.

The fractal structure of branched polymers embedded in $\mathbb{R}^D$ is described by the extrinsic Hausdorff dimension D_H. D_H can be computed from the scaling of the embedded two-point function. The path between two marked points x and y is an embedded random walk at which at each point branched polymers can grow out. The integration over the embedding field is the same

as for the ordinary random walk in D dimensions. The branching at the vertices corresponds to a renormalization of $e^{-\mu}$. From this renormalization D_H can be computed with the result $D_H = \frac{2}{\gamma} = \frac{2m}{m-1}$ [21]. Thus $D_H = 4$ for ordinary branched polymers and $D_H \to 2$ for $m \to \infty$.

2.2 Matrix models

Major advances in the theory of two-dimensional quantum gravity have been made when it was realized, that the theory is integrable in the discrete formulation, even for some types of matter coupled to the surfaces [56, 70, 57, 71, 72, 58, 73].

In the development of these results the formulation of dynamical triangulation in terms of matrix models has been of some importance. The Hermitean one-matrix model has a direct interpretation in terms of randomly triangulated surfaces. This is demonstrated in the beginning of this section. Also the Hartle-Hawking wavefunctionals have a direct natural formulation via matrix models. The Dyson-Schwinger equations for these models allow the treatment of the theory by methods of complex analysis. These so-called loop equations can alternatively be derived using only combinatorial methods [62]. The explicit solution of the loop equations provides deep insight into the properties of the theory and allows the computation of the scaling limit of the Hartle-Hawking wavefunctionals.

We cannot review the vast literature on numerical simulations in quantum gravity in this work. However, let us remark that all numerical simulations of two-dimensional dynamical triangulation are consistent with the scaling hypotheses, with universality, and – most remarkably – with the continuum theory of two-dimensional quantum gravity.

2.2.1 Dynamical triangulation by matrix models

The discretized partition function of two-dimensional quantum gravity can be represented as an integral over a Hermitean $N \times N$-matrix ϕ. Consider the Gaussian integral

$$\int d\phi \; e^{-\frac{1}{2}\mathrm{Tr}\phi^2} \frac{1}{K!} \left(\frac{1}{3}\mathrm{Tr}\phi^3\right)^K, \tag{2.54}$$

where the measure $d\phi$ is defined as

$$d\phi = \prod_{i \le j} d\Re\phi_{ij} \prod_{i<j} d\Im\phi_{ij}. \tag{2.55}$$

The propagator of ϕ is given by

$$\langle \phi_{ij}\phi_{i'j'}\rangle \equiv \frac{1}{Z(N,0)}\int d\phi\ e^{-\frac{1}{2}\mathrm{Tr}\phi^2}\phi_{ij}\phi_{i'j'} = \delta_{ij'}\delta_{ji'}, \tag{2.56}$$

with $Z(N,0) = \int d\phi\ e^{-\frac{1}{2}\mathrm{Tr}\phi^2}$. Diagrammatically, the propagator can be represented as a double line where the two lines are oriented in opposite directions. The integral (2.54) is performed by doing all possible contractions of the K $\mathrm{Tr}\phi^3$ vertices. The dual picture of this corresponds to gluing K triangles together to form all possible closed and not necessarily connected surfaces of arbitrary topology. The contribution from a particular graph forming a closed surface is N^{N_v}, since each vertex contributes with a factor N. If we make the substitution

$$\mathrm{Tr}\phi^3 \to \frac{g}{\sqrt{N}}\mathrm{Tr}\phi^3, \tag{2.57}$$

each closed surface with Euler characteristic χ contributes with $g^K N^{N_v - K/2} = g^K N^\chi$. We get the sum over arbitrary closed surfaces with any number of triangles if we sum (2.54) over K:

$$Z(N,g) = \int d\phi\ e^{-\frac{1}{2}\mathrm{Tr}\phi^2 + \frac{g}{3\sqrt{N}}\mathrm{Tr}\phi^3}. \tag{2.58}$$

Taking the logarithm of this projects on the connected surfaces only. Thus we see that

$$Z(\mu, G) = \log\frac{Z(N,g)}{Z(N,0)}, \tag{2.59}$$

with

$$\frac{1}{G} = \log N, \text{ and } \mu = -\log g \tag{2.60}$$

is a formal definition of the discretized partition function of two-dimensional quantum gravity, including the sum over topologies. Here G is the gravitational coupling constant. The matrix integral (2.58) is of course not convergent, but it has been suggested, that a closed form like (2.58) might define a non-perturbative definition of the sum over topologies after analytic continuation. In general however, the integral will be complex [74, 75] and the problem of summing over topologies has not yet been solved.

Equation (2.58) admits a $1/N^2$-expansion which is identical to an expansion over topologies. Here we will restrict ourselves to spherical topology

which corresponds to the leading order in $1/N^2$. It is convenient to generalize the matrix integral (2.58) to

$$Z(N, g_i) = \int d\phi \; e^{-N\mathrm{Tr}V(\phi)}, \tag{2.61}$$

$$V(\phi) = \sum_{n=1}^{\infty} \frac{g_n}{n}\phi^n, \tag{2.62}$$

where we have performed the rescaling $\phi \to \sqrt{N}\phi$. (2.61) is interpreted as a perturbative expansion around a Gaussian integral. Thus the Gaussian coupling constant is bigger than zero, while the others are chosen smaller than zero. The expectation value of some observable $\mathcal{O}(\phi)$ is defined as

$$\langle \mathcal{O}(\phi) \rangle = \frac{1}{Z(N, g_i)} \int d\phi \; e^{-N\mathrm{Tr}V(\phi)} \mathcal{O}(\phi). \tag{2.63}$$

Differentiating the logarithm of (2.61) with respect to the coupling constants g_n defines expectation values of observables like $\mathrm{Tr}\phi^{k_1} \cdots \mathrm{Tr}\phi^{k_b}$. They are interpreted as the sum over all surfaces which have b polygons with k_i links as their boundary. The generating function of the connected expectation values of these observables is given by

$$\begin{aligned} W(z_1, \ldots, z_b) &= N^{b-2} \sum_{k_1, \ldots, k_b = 0}^{\infty} \frac{\left\langle \mathrm{Tr}\phi^{k_1} \cdots \mathrm{Tr}\phi^{k_b} \right\rangle_{\mathrm{conn}}}{z_1^{k_1+1} \cdots z_b^{k_b+1}} \\ &= N^{b-2} \left\langle \mathrm{Tr}\frac{1}{z_1 - \phi} \cdots \mathrm{Tr}\frac{1}{z_b - \phi} \right\rangle_{\mathrm{conn}}. \end{aligned} \tag{2.64}$$

In the large-N limit this is the generating function for the discretized Hartle-Hawking wavefunctionals, whose continuum counterparts have been defined in section 1.1.2. In fact, already the one-loop correlator $W(z)$ contains all necessary information, since the higher correlators can be computed from it by differentiation:

$$W(z_1, \ldots, z_b) = \frac{d}{dV(z_b)} \frac{d}{dV(z_{b-1})} \cdots \frac{d}{dV(z_2)} W(z_1), \tag{2.65}$$

where the loop insertion operator $\frac{d}{dV(z)}$ is defined as

$$\frac{d}{dV(z)} \equiv -\sum_{n=1}^{\infty} \frac{n}{z^{n+1}} \frac{\partial}{\partial g_n}. \tag{2.66}$$

Let us define the density $\rho(\lambda)$ of eigenvalues of the matrix integral (2.61) by

$$\rho(\lambda) = \left\langle \sum_{i=1}^{N} \delta(\lambda_i - \lambda) \right\rangle, \tag{2.67}$$

where the λ_i are the N eigenvalues of the matrix ϕ. Then the one-loop correlator $W(z)$ can be written as:

$$W(z) = \int_{-\infty}^{\infty} d\lambda \, \frac{\rho(\lambda)}{z-\lambda}. \tag{2.68}$$

For polynomial potentials the support of $\rho(\lambda)$ is confined to one or several intervals on the real axes in the limit $N \to \infty$. The solution in the case of several cuts has been given recently [76]. Here we will only deal with a single cut $[a, b]$ on the real axes. Then $W(z)$ will be analytic in the complex plane with a cut at the support $[a, b]$ of $\rho(\lambda)$. It follows that

$$2\pi i \, \rho(\lambda) = \lim_{\epsilon \to 0} \left(W(\lambda + i\epsilon) - W(\lambda - i\epsilon) \right). \tag{2.69}$$

2.2.2 The loop equations

The matrix model (2.61) can be solved by many methods. It is most conveniently and systematically done by using loop equations, the Dyson-Schwinger equations for the matrix models. Let us consider the transformation

$$\phi \to \phi + \epsilon \frac{1}{z-\phi}, \tag{2.70}$$

which makes sense if z is not an eigenvalue of ϕ and real, so that the new matrix remains Hermitean. Then the measure and the action transform as

$$d\phi \;\; \to \;\; \left(1 + \epsilon \mathrm{Tr} \frac{1}{z-\phi} \mathrm{Tr} \frac{1}{z-\phi} \right) d\phi, \tag{2.71}$$

$$\mathrm{Tr} V(\phi) \;\; \to \;\; \mathrm{Tr} V(\phi) + \epsilon \mathrm{Tr} \frac{V'(\phi)}{z-\phi}, \tag{2.72}$$

to first order in ϵ. Since the matrix integral (2.61) is invariant under such a definition of the integration variable we get:

$$\int d\phi \; e^{-N \mathrm{Tr} V(\phi)} \left\{ \left(\mathrm{Tr} \frac{1}{z-\phi} \right)^2 - N \mathrm{Tr} \frac{V'(\phi)}{z-\phi} \right\} = 0. \tag{2.73}$$

The second term can be rewritten as a contour integral involving the one-loop correlator by using the eigenvalue density $\rho(\lambda)$, while the first term is related to the two-loop correlator. This leads to the standard form of the loop equations [77]:

$$\oint_C \frac{d\omega}{2\pi i} \, \frac{V'(\omega)}{z-\omega} W(\omega) = W(z)^2 + \frac{1}{N^2} W(z, z). \tag{2.74}$$

The contour $\mathcal{C}$ goes around the cut $[a, b]$ but does not enclose z. The loop equation can be solved systematically to all orders in $1/N^2$, that means for all genera of the random surfaces [78, 79]. For spherical topology, that means in the limit $N \to \infty$, (2.74) simplifies to

$$\oint_{\mathcal{C}} \frac{d\omega}{2\pi i} \frac{V'(\omega)}{z-\omega} W_0(\omega) = W_0(z)^2, \tag{2.75}$$

where the subscript 0 denotes the genus of the surfaces. By the definition we know that $W_0(z) = \frac{1}{z} + O(z^{-2})$. Therefore (2.75) can be solved by a deformation of the contour $\mathcal{C}$ to infinity. For a polynomial action V of degree n we get:

$$\begin{aligned} W_0(z) &= \frac{1}{2}\left(V'(z) - \sqrt{(z-a)(z-b)} \sum_{k=1}^{\infty} M_k[a, b, g_i](z-b)^{k-1} \right) \\ &= \oint_{\mathcal{C}} \frac{d\omega}{2\pi i} \frac{V'(\omega)}{z-\omega} \frac{\sqrt{(z-a)(z-b)}}{\sqrt{(\omega-a)(\omega-b)}}, \end{aligned} \tag{2.76}$$

where the moments M_k are defined as

$$M_k[a, b, g_i] = \oint_{\mathcal{C}} \frac{d\omega}{2\pi i} \frac{V'(\omega)}{(\omega-a)^{\frac{1}{2}}(\omega-b)^{k+\frac{1}{2}}}, \tag{2.77}$$

and vanish for $k > n-1$. The endpoints a and b of the cut are self-consistently determined by the equations

$$M_{-1}[a, b, g_i] = 2, \tag{2.78}$$

$$M_0[a, b, g_i] = 0, \tag{2.79}$$

which are a consequence of (2.76) together with $W_0(z) = \frac{1}{z} + O(z^{-2})$.

For the matrix model (2.58) which directly corresponds to triangulated surfaces, the solution for $W_0(z)$ is:

$$\begin{aligned} W_0(z) &= \frac{1}{2}V'(z) + f(\mu, z), \\ f(\mu, z) &= \frac{g}{2}(z-c)\sqrt{(z-a)(z-b)}, \end{aligned} \tag{2.80}$$

where $g = e^{-\mu}$ and $c = \frac{1}{g} - \frac{a+b}{2}$.

In principle one could compute all higher loop correlators from (2.76) by applying the loop insertion operator (2.66), which gives a complete solution of two-dimensional dynamical triangulation for spherical topology.

In fact, a closed expression for the b-loop correlators can be obtained if the small scale details of the theory are adjusted. Instead of the Hermitean matrix model one considers the complex matrix model with a potential

$$\mathrm{Tr}V(\phi^+\phi) = \sum_{n=1}^{\infty} \frac{g_n}{n} \mathrm{Tr}(\phi^+\phi)^n, \tag{2.81}$$

and an integration measure

$$d\phi = \prod_{i,j=1}^{N} d\Re\phi_{ij} d\Im\phi_{ij}. \tag{2.82}$$

The one-loop correlator is now defined as

$$W(z) = \frac{1}{N} \sum_{n=0}^{\infty} \frac{\langle \mathrm{Tr}(\phi^+\phi)^n \rangle}{z^{2n+1}}, \tag{2.83}$$

higher loop correlators are defined analogously. The term $\mathrm{Tr}(\phi^+\phi)^n$ can be interpreted as a $2n$-sided polygon whose links are alternately coloured black and white. These polygons can be glued together as in the Hermitean matrix model with the additional constraint that black links have to be glued to white links [80]. Such short distance details about the gluing should be unimportant in the continuum limit.

The loop equations for this model have been derived in [81]. Because of the symmetry $\phi \to -\phi$ we have $a = -b$ for the cut of the one-loop correlator. The solution for spherical topology is given by [81]:

$$\begin{aligned} W_0(z) &= \frac{1}{2}\left(V'(z) - M(z)\sqrt{z^2 - b^2}\right), \\ M(z) &= \oint_{\mathcal{C}_\infty} \frac{d\omega}{4\pi i} \frac{\omega V'(\omega)}{(\omega^2 - z^2)\sqrt{\omega^2 - b^2}} = \sum_{k=1}^{\infty} M_k[b, g_i](z^2 - b^2)^{k-1}, \\ M_k[b, g_i] &= \oint_{\mathcal{C}} \frac{d\omega}{4\pi i} \frac{\omega V'(\omega)}{(\omega^2 - b^2)^{k+\frac{1}{2}}}. \end{aligned} \tag{2.84}$$

Here $\mathcal{C}_\infty$ is a contour around the cut pushed to infinity. The position of the cut is determined by the equation

$$M_0[b, g_i] = 2. \tag{2.85}$$

The higher loop correlators are given by the following expressions [81]:

$$\begin{aligned} W_0(z_1,z_2) &= \frac{1}{4(z_1^2-z_2^2)^2}\left(z_2^2\sqrt{\frac{z_1^2-b^2}{z_2^2-b^2}}+z_1^2\sqrt{\frac{z_2^2-b^2}{z_1^2-b^2}}-2z_1z_2\right), \\ W_0(z_1,z_2,z_3) &= \frac{b^4}{16M_1}\frac{1}{\sqrt{(z_1^2-b^2)(z_2^2-b^2)(z_3^2-b^2)}}, \\ W_0(z_1,\ldots,z_b) &= \left(\frac{1}{M_1}\frac{d}{db^2}\right)^{b-3}\frac{1}{2b^2M_1}\prod_{k=1}^{b}\frac{b^2}{2(z_k^2-b^2)^{3/2}}. \end{aligned} \tag{2.86}$$

In these formulas all dependence on the coupling constants is hidden in M_1 and b. Similar and only slightly more complicated statements are valid also for the Hermitean matrix model.

2.2.3 Scaling limit and computation of γ

It has been discussed in section 2.1.3 that the ensemble of triangulated random surfaces has a critical point μ_c. The continuum limit with a renormalized cosmological constant Λ is approached by $\mu \to \mu_c$ with $\mu - \mu_c = \Lambda a^2$, where a is the lattice spacing. In terms of the coupling constant g for the model (2.58) this relation is $g_c - g \sim g_c \Lambda a^2$, close to the critical point. For the generalized matrix model (2.61) with n coupling constants we expect no qualitative changes except that the theory will be critical on a $(n-1)$-dimensional hypersurface. This hypersurface is identified by

$$M_1[b(g_i), g_i] = 0, \tag{2.87}$$

since the b-loop correlators $W_0(z_1,\ldots,z_b)$ are exactly divergent for $M_1 = 0$. Let us denote a point in the critical hypersurface by $g_{c,i}, i = 1,\ldots,n$ and the corresponding endpoint of the eigenvalue distribution by b_c. If we move slightly away from the critical surface

$$g_i = g_{c,i}(1 - \Lambda a^2) = g_{c,i} + \delta g_i, \tag{2.88}$$

there will be a corresponding change $b_c^2 \to b_c^2 + \delta b^2$ which can be computed from (2.85) to be

$$(\delta b^2)^2 = -\frac{16}{3M_2[b_c, g_{c,i}]}\Lambda a^2 \sim \delta g_i. \tag{2.89}$$

We rescale the cosmological constant such that $b^2 = b_c^2 - a\sqrt{\Lambda}$. Because z_i appears in (2.86) always in the combination $(z_i^2 - b^2)$ it is natural to introduce a scaling of z_i by

$$z_i^2 = b_c^2 + aZ_i. \tag{2.90}$$

Since $W_0(z_1, \ldots, z_b)$ is the generating function for b-loop correlators, we can compute the transition amplitude for b one-dimensional universes of lengths $n_1, \ldots, n_b$ from it by a multiple contour integration. The physical lengths of the loops will be $L_i = n_i a$. We see that in the continuum limit the number of links n_i on the boundaries has to go to infinity as $a \to 0$ if the loops shall survive. By following this procedure one gets the generating functional for macroscopic b-loop amplitudes [81]:

$$\begin{aligned} W_0(z_1, \ldots, z_b) &\sim a^{5-\frac{7b}{2}} W_0(\Lambda, L_1, \ldots, L_b), \\ W_0(\Lambda, Z_1, \ldots, Z_b) &= \left(-\frac{d}{d\Lambda}\right)^{b-3} \frac{1}{\sqrt{\Lambda}} \prod_{k=1}^{b} (Z_k + \sqrt{\Lambda})^{-\frac{3}{2}}, \end{aligned} \tag{2.91}$$

for $b \geq 3$. The inverse Laplace transform of $W_0(\Lambda, Z_1, \ldots, Z_b)$ in the variables Z_i gives the Hartle-Hawking wavefunctionals of two-dimensional quantum gravity. It can be computed from (2.91) to be:

$$W_0(\Lambda, L_1, \ldots, L_b) = \left(-\frac{d}{d\Lambda}\right)^{b-3} \frac{1}{\sqrt{\Lambda}} \sqrt{L_1 \cdots L_b}\; e^{-\sqrt{\Lambda}(L_1 + \ldots + L_b)}. \tag{2.92}$$

Since the b-point function in two-dimensional quantum gravity should scale as $\Lambda^{2-b-\gamma}$, we can directly read off from this expression that

$$\gamma = -\frac{1}{2} \tag{2.93}$$

for pure gravity, in agreement with the KPZ-formula (1.14). These calculations can be generalized to arbitrary topology [77, 78, 82, 79] and to models with matter coupled to gravity [83, 84, 62] by considering multicritical models which are obtained by a fine-tuning of the critical coupling constants such that higher moments vanish as well as M_1.

It is a major result that all values of the susceptibility exponent γ computed by these or other methods in the model of dynamically triangulated quantum gravity agree with the continuum formula (1.14) which has been derived in section 1.2.3. In fact, all calculations which can be done by dynamical triangulation and in the continuum approach have so far yielded the same results. Therefore we identify both theories as the theory of two-dimensional quantum gravity. Objects like the Hartle-Hawking wavefunctionals and other correlation functions can much easier be obtained in the discretized approach whose scaling limit yields the theory of two-dimensional quantum gravity.

2.3 The fractal structure of pure gravity

Another major advance in two-dimensional quantum gravity was made by the explicit computation of the two-point function for pure quantum gravity by constructing a transfer matrix [22] and later by using a peeling decomposition [23]. The two-point function is a natural object since it provides all details about the scaling properties of two-dimensional quantum gravity. The astonishing result of these investigations is that the intrinsic Hausdorff dimension of two-dimensional quantum gravity is *four*. That means that even the dimensionality of spacetime is a dynamical quantity.

2.3.1 The geodesic two-loop function

Let us define with $\mathcal{T}(l_1, l_2, r)$ the class of triangulations with an entrance loop l_1 with one marked link and an exit loop l_2, separated a geodesic distance r. We will also denote the number of links of l_1 and l_2 with the same symbols. Then the geodesic two-loop function is defined as

$$G(\mu, r; l_1, l_2) = \sum_{T \in \mathcal{T}(l_1, l_2, r)} e^{-\mu N_t}. \tag{2.94}$$

For $r = 0$ we have the initial condition:

$$G(\mu, 0; l_1, l_2) = \delta_{l_1, l_2}. \tag{2.95}$$

We introduce the generating function for $G(\mu, r; l_1, l_2)$ by

$$G(\mu, r; z_1, z_2) = \sum_{l_1, l_2 = 1}^{\infty} z_1^{-(l_1+1)} z_2^{-(l_2+1)} G(\mu, r; l_1, l_2), \tag{2.96}$$

with the initial condition

$$G(\mu, 0; z_1, z_2) = \frac{1}{z_1 z_2} \frac{1}{z_1 z_2 - 1}. \tag{2.97}$$

By a two-fold contour integration the geodesic two-loop function can be reconstructed:

$$G(\mu, r; l_1, l_2) = \oint_{\mathcal{C}_1} \frac{dz_1}{2\pi i}\, z_1^{l_1} \oint_{\mathcal{C}_2} \frac{dz_2}{2\pi i}\, z_2^{l_2}\, G(\mu, r; z_1, z_2). \tag{2.98}$$

It is an important observation that the two-point function $G(\mu, r)$ can be obtained from the geodesic two-loop function. Consider the two-loop function with an entry loop of length $l_1 = 1$ which is equivalent to a marked link,

and an exit loop of arbitrary length l_2 separated a geodesic distance r. We can close this surface by gluing a disc to the boundary l_2. The amplitude $W_0(l_2)$ of the disc can be computed from the one-loop correlator $W_0(z)$ by a contour integration. An additional factor of l_2 arises because we have to mark one of the l_2 links of the exit loop. Thus we get:

$$\begin{aligned} G(\mu, r) &= \sum_{l_2=1}^{\infty} G(\mu, r; 1, l_2)\ l_2 W_0(l_2) \\ &= \oint_{\mathcal{C}} \frac{d\omega}{2\pi i} \frac{1}{\omega} \left[z^2 G(\mu, r; z, \frac{1}{\omega}) \right] \left[-\frac{\partial}{\partial \omega} \omega W_0(\omega) \right] \Big|_{z=\infty}, \end{aligned} \tag{2.99}$$

where $\mathcal{C}$ is a contour around zero. Thus we see that all information about the scaling of the theory can be obtained if the generating function of geodesic two-loop amplitudes can be computed.

By a step-by-step decomposition of triangulations in $\mathcal{T}(l_1, l_2, r)$ a differential equation for $G(\mu, r; z, \omega)$ can be obtained . It is intuitive that any triangulation in $\mathcal{T}(l_1, l_2, r)$ can be decomposed into r rings of thickness one [22]. This leads to a transfer matrix formalism for the two-point function. Alternatively, one can decompose the triangulations by a peeling procedure [23]. The differential equation one obtains is not exact but should be valid close to the critical point of the theory:

$$\frac{\partial}{\partial r} G(\mu, r; z, \omega) = -2 \frac{\partial}{\partial z} \left[f(\mu, z) G(\mu, r; z, \omega) \right], \tag{2.100}$$

where $f(\mu, z)$ is given by (2.80) for triangulated surfaces.

2.3.2 Scaling of the two-point function

The differential equation (2.100) can be solved by the method of characteristic equations. The result is:

$$G(\mu, r; z, \omega) = \frac{f(\mu, \hat{z})}{f(\mu, z)} \frac{\hat{z}\omega}{\hat{z}\omega - 1}, \tag{2.101}$$

where $\hat{z}$ is the solution to the characteristic equation

$$\frac{d\hat{z}(z, r)}{dr} = 2 f(\mu, \hat{z}), \tag{2.102}$$

given by

$$\frac{1}{\hat{z}(z, r)} = \frac{1}{c} - \frac{\delta_1}{c} \frac{1}{\sinh^2 \left(-\delta_0 r + \sinh^{-1} \sqrt{\frac{\delta_1}{1 - c/z} - \delta_2} \right) + \delta_2}. \tag{2.103}$$

Here the positive constants δ_i scale as

$$\delta_0 = \frac{g}{2}\sqrt{(c-a)(c-b)} = O((\mu-\mu_c)^{\frac{1}{4}}), \tag{2.104}$$

$$\delta_1 = \frac{(c-a)(c-b)}{c(b-a)} = O((\mu-\mu_c)^{\frac{1}{2}}), \tag{2.105}$$

$$\delta_2 = -\frac{a(c-b)}{c(b-a)} = O((\mu-\mu_c)^{\frac{1}{2}}). \tag{2.106}$$

Now the two-point function $G(\mu, r)$ can be computed by inserting (2.101) into (2.99). Close to the critical point one obtains [85]:

$$G(\mu, r) = \text{const} \times \delta_0\delta_1 \frac{\cosh(\delta_0 r)}{\sinh^3(\delta_0 r)}(1 + O(\delta_0)). \tag{2.107}$$

From (2.104) we get:

$$G(\mu, r) = \text{const} \times (\mu-\mu_c)^{\frac{3}{4}} \frac{\cosh\left(c(\mu-\mu_c)^{\frac{1}{4}} r\right)}{\sinh^3\left(c(\mu-\mu_c)^{\frac{1}{4}} r\right)}, \tag{2.108}$$

where $c = \sqrt{6}e^{-\mu_c}$ is a nonuniversal constant. To read off the critical behaviour of the theory we only have to analyze the asymptotic behaviour of the two-point function for large and for small distances. We see that:

- $G(\mu, r)$ falls off exponentially as $e^{-2c(\mu-\mu_c)^{\frac{1}{4}} r}$, for $r \to \infty$. Thus the susceptibility exponent equals $\nu = \frac{1}{4}$ and the intrinsic Hausdorff dimension d_H of pure two-dimensional quantum gravity equals *four*.

- For $1 \ll r \ll (\mu-\mu_c)^{-\frac{1}{4}}$ the two-point function behaves like r^{-3}, that means the anomalous scaling dimension is $\eta = 4$. For ordinary statistical systems η is smaller than two. It is remarkable, that the critical exponents of the theory still satisfy Fisher's scaling relation $\gamma = \nu(2-\eta)$.

- Any geodesic two-loop function $G(\mu, r; l_1, l_2)$, which can be computed from (2.98) and (2.101) fulfills the same scaling relations as the two-point function, provided that l_1 and l_2 stay finite in the limit $\mu \to \mu_c$.

- The continuum limit of the two-point function is given by

$$G(\Lambda, R) = C\Lambda^{\frac{3}{4}} \frac{\cosh\left(c\Lambda^{\frac{1}{4}} R\right)}{\sinh^3\left(c\Lambda^{\frac{1}{4}} R\right)}, \tag{2.109}$$

  where C is a constant.

2.4 Conclusion

In this chapter we have demonstrated, how to use discretized systems to describe two-dimensional quantum gravity. In two dimensions, the continuum theory of quantum gravity and the theory of dynamical triangulations agree with each other in that region, where the continuum theory can be evaluated, that means for matter with central charge $D \leq 1$ coupled to gravity. The continuum limit of dynamical triangulations in two dimensions can therefore be identified with the continuum theory.

In fact, the discretized approach is more powerful than the continuum approach in the sense that natural observables like the Hartle-Hawking wavefunctionals can be computed. Also the two-point function for pure quantum gravity can be obtained. This object is of central interest when questions about the scaling of the theory are to be addressed.

On the technical side we have reviewed matrix models, which have developed a status of interest on their own beyond dynamical triangulation in string theory and in condensed matter theory.

Since there is at present no successful way to find a continuum theory of quantum gravity in higher dimensions than two, it is natural to ask, whether such a theory could be defined as the scaling limit of the corresponding higher dimensional discretized theory. It turns out that dynamical triangulation can be defined in three and four dimensions and that these theories have a phase transition similar to the two-dimensional case. While there has been some analytical progress in the understanding of entropy bounds on the number of triangulations in higher dimensions [86] most work in this field is numerical. Questions about the nature of the phase transition in four dimensions are not settled yet and exciting research remains to be done.

Chapter 3

The failure of quantum Regge calculus

In the continuum path integral formulation of quantum gravity we are instructed to compute the integral

$$\int \mathcal{D}[g_{\mu\nu}] e^{-S_{EH}(g)} \equiv \int \frac{\mathcal{D}g_{\mu\nu}}{\mathrm{vol}(\mathrm{Diff})} e^{-S_{EH}(g)} \tag{3.1}$$

over diffeomorphism classes of metrics weighted with the exponential of the Einstein-Hilbert action, as has been discussed in chapter 1. Two different discretization schemes have been suggested. One, dynamical triangulation, provides a regularization of the functional integral (3.1). An explicit cutoff is introduced and one sums over all (abstract) triangulations with equilateral d-simplices of a d-dimensional manifold M, compare chapter 2.

A different attempt to discretize the functional integral (3.1), called quantum Regge calculus (QRC), has been suggested, see [87, 88, 89] for reviews and references. This formalism is closely related to the classical coordinate independent Regge discretization of general relativity: One *fixes* a (suitably chosen) triangulation, while the dynamical degrees of freedom are the L link lengths. Thus the functional integration in (3.1) is replaced by

$$\int d\mu(l_1, \ldots, l_L) \equiv \int_0^\infty \prod_{i=1}^{L} dl_i \; J(l_1, \ldots, l_L) \; \delta(\Delta), \tag{3.2}$$

where $J(l_1, \ldots, l_L)$ is the Jacobian of the transformation $dg_{\mu\nu}(\xi) \to dl_i$. The integral in (3.2) is over all link length assignments consistent with the triangle inequalities, as denoted by the delta function $\delta(\Delta)$. While it provides a discretization of the integration in (3.1), this replacement does not provide a regularization of (3.1). Contrary to dynamical triangulation, no cutoff has

to be introduced [90]. One can of course choose to work with a cutoff which has to be taken to zero at the end of the computations. This will not make any difference in the argumentation below.

The Jacobian in (3.2) is very complicated and its form is presently unknown. For analytical and numerical investigations it is usually replaced by local measures of the form $\prod_{i=1}^{L} f(l_i)$.

If a sensible continuum limit of this theory existed, it would be approached by taking the number L of links in the triangulation to infinity at a critical point of the statistical ensemble defined by (3.2) and by the discretized Einstein Hilbert action. For this limit to make sense, the resulting continuum theory should not depend on details of the discretized theory like the chosen fixed triangulation or the local measure $f(l_i)$.

It is one central objective of this chapter to show analytically that such a continuum limit of quantum Regge calculus in its present formulation cannot be defined in any dimension $d > 1$.

Such a result has been indicated by simulations in two-dimensional quantum Regge calculus [91, 92, 93]. These simulations revealed that the KPZ-exponents (1.14) for the susceptibility of conformal matter coupled to continuum quantum gravity could not be obtained. This has often been called the failure of quantum Regge calculus. Most discussion about this failure to reproduce continuum results in two dimensions has been centered around the choice of the measure.

In the first part of this chapter some notation is introduced. In the second part we discuss the measures which have been suggested and used in the context of quantum Regge calculus. In the third part we show that in two dimensions all discussed measures for Regge calculus fail to reproduce continuum results. Not even the concept of length can be defined [2]. In the conclusion we discuss the situation in higher dimensions. We give a short review of an alternative approach, in which the continuum functional integral over metrics is restricted to piecewise linear metrics [94].

This chapter is partly based on work presented in [2].

3.1 Formulation of quantum Regge calculus

Let us fix the connectivity of a triangulation with V vertices[1], T triangles and L links of lengths $l_1, \ldots, l_L$ of a two-dimensional closed manifold M with Euler-characteristic χ. The number of links can also be expressed as

[1] We denote the volume of manifolds with the same symbol V. No confusion should arise.

$L = 3V - 3\chi$. We view the interior of the simplices as flat. As explained in section 2.1.1 the curvature of this piecewise linear space is concentrated at the vertices and each assignment of link lengths uniquely determines a metric on M. For a triangle this metric can be written as

$$g_{\mu\nu} = \begin{pmatrix} x_1 & \frac{1}{2}(x_1 + x_2 - x_3) \\ \frac{1}{2}(x_1 + x_2 - x_3) & x_2 \end{pmatrix}, \tag{3.3}$$

where the squared edge lengths are defined as $x_i = l_i^2,\ i = 1, \ldots, L$. The area A of a single triangle can be expressed as

$$A = \int d^2\xi \sqrt{g(\xi)} = \frac{1}{2}\sqrt{g} = \frac{1}{2}\sqrt{x_1 x_2 - \frac{1}{4}(x_1 + x_2 - x_3)^2}. \tag{3.4}$$

Since in quantum Regge calculus the link lengths are the dynamical variables one attempts to replace the functional integration over equivalence classes of metrics by an integration over all link length assignments which correspond to different geometries. Clearly, this replacement

$$\int \mathcal{D}[g_{\mu\nu}] = \int \frac{\mathcal{D}g_{\mu\nu}}{\mathrm{Vol(Diff)}} \to \int \prod_{i=1}^{L} dl_i \; J(l_1, \ldots, l_L)\delta(\Delta) \equiv \int d\mu(l_1, \ldots, l_L) \tag{3.5}$$

involves a highly non-trivial Jacobian $J(l_1, \ldots, l_L)$. The integration is over all link lengths compatible with the triangle inequalities. But not all assignments of link lengths define independent geometries as can be seen in the flat case. All vertices can be moved around in the plane, changing the link lengths without changing the flat geometry. The Jacobian has to be such that this additional invariance is divided out of the integral.

Thus, we obtain an $L = 3V - 3\chi$-dimensional subspace of the infinite dimensional space of equivalence classes of metrics. Quantum Regge calculus replaces the functional integration in (1.1) by an integral over this finite dimensional subspace. It is hoped that in the limit $L \to \infty$ expectation values of observables converge in some suitable way to their continuum values.

3.2 Regge integration measures

Presently, the form of the Jacobian in (3.5) is not known. However, by appealing to universality one might assume that the choice of this measure is not very important. A large number of measures have been suggested and tried out in numerical experiments to study whether quantum Regge calculus in its present form agrees with the results of continuum calculations

and dynamical triangulations in two dimensions. The result of these investigations is negative. In this section we will define the most important and common of these measures and study some of their properties.

3.2.1 DeWitt-like measures

One possible way to construct a measure for quantum Regge calculus in terms of the link lengths is to repeat the construction in section 1.2.1 with the constraint that the deformations of the metric are allowed by quantum Regge calculus. For an alternative discussion see [89] where the corresponding metric is called Lund-Regge metric, refering to an unpublished work by Lund and Regge. However, since the starting point of our discussion is the DeWitt metric we rather call the measure a DeWitt-like measure. Note however that it is not the DeWitt measure restricted to piecewise linear metrics. This will be given in section 3.4.2.

For a single triangle the variation of the metric (3.3) in terms of the δx_i is given by

$$\delta g_{\mu\nu} = \begin{pmatrix} \delta x_1 & \frac{1}{2}(\delta x_1 + \delta x_2 - \delta x_3) \\ \frac{1}{2}(\delta x_1 + \delta x_2 - \delta x_3) & \delta x_2 \end{pmatrix}. \tag{3.6}$$

Using this and the DeWitt metric, the scalar product $\langle \delta g, \delta g \rangle_T$, which has been defined in section 1.2.1 can be computed for a single triangle with area A. It turns out that it simplifies considerably when we use the canonical value -2 for the parameter C in the DeWitt metric:

$$\begin{aligned} \langle \delta g, \delta g \rangle_T &= \int d^2\xi \sqrt{g(\xi)}\, \delta g_{\mu\nu}(\xi) G^{\mu\nu,\alpha\beta} \delta g_{\alpha\beta}(\xi) \\ &= \frac{A}{2} \left(2\delta g_{\mu\nu} g^{\mu\alpha} g^{\nu\beta} \delta g_{\alpha\beta} + C \delta g_{\mu\nu} g^{\mu\nu} g^{\alpha\beta} \delta g_{\alpha\beta} \right) \\ &= -2A \det(\delta g_{\mu\nu}) \det(g^{\mu\nu}), \text{ for } C = -2. \end{aligned} \tag{3.7}$$

Using (3.3) and (3.6) leads to

$$\langle \delta g, \delta g \rangle_T = [\delta x_1, \delta x_2, \delta x_3] \frac{1}{16A} \begin{bmatrix} 1 & -1 & -1 \\ -1 & 1 & -1 \\ -1 & -1 & 1 \end{bmatrix} \begin{bmatrix} \delta x_1 \\ \delta x_2 \\ \delta x_3 \end{bmatrix}. \tag{3.8}$$

For a general two-dimensional triangulation this line element is given by the

sum of (3.8) over all triangles:

$$\begin{aligned}\langle \delta g, \delta g\rangle_T &= \sum_t \int d^2\xi \sqrt{g^t(\xi)}\, \delta g^t_{\mu\nu}(\xi)(G^t)^{\mu\nu\alpha\beta}\delta g^t_{\alpha\beta}(\xi) \\ &= [\,\delta x_1, \ldots, \delta x_L\,]\, M \begin{bmatrix} \delta x_1 \\ \vdots \\ \delta x_L \end{bmatrix}. \end{aligned} \tag{3.9}$$

Here M is an $L \times L$ matrix. M_{ij} is the element corresponding to the links l_i and l_j. If both l_i and l_j belong to the same triangle t with area A_t the corresponding non-diagonal entry is

$$M_{ij} = M_{ji} = -\frac{1}{A_t}. \tag{3.10}$$

All other off-diagonal entries are zero. It follows that M has four nonvanishing off-diagonal entries in each row. Diagonal entries are given by

$$M_{ii} = \frac{1}{A_{t_1}} + \frac{1}{A_{t_2}}, \tag{3.11}$$

where t_1 and t_2 are the two triangles which contain l_i. From the i'th row of M the product $(A_{t_1}A_{t_2})^{-1}$ can be factorized. Since each triangle has three sides, this amounts to factorizing $\prod_{k=1}^{T} A_k^{-3}$ from the determinant of M:

$$\begin{aligned}\det M &= \prod_{k=1}^{T} A_k^{-3} \begin{vmatrix} A_1+A_2 & -A_2 & -A_2 & -A_1 & -A_1 & 0 & \ldots \\ & & & \ldots & & & \end{vmatrix} \\ &=: P(A_1, \ldots, A_T) \prod_{k=1}^{T} A_k^{-3}. \end{aligned} \tag{3.12}$$

$P(A_1, \ldots, A_T)$ is a polynomial in the areas of the triangles which vanishes, whenever two adjacent triangle areas vanish. It follows directly that P is a highly nonlocal function of the areas, since each monomial of P must contain at least half of the triangles.

Therefore the DeWitt-like integration measure for quantum Regge calculus is given by

$$d\mu(l_1, \ldots, l_L) = \text{const} \times \frac{\sqrt{P(A_1, \ldots, A_T)}}{\prod_{k=1}^{T} A_k^{3/2}} \prod_{j=1}^{L} l_j dl_j\, \delta(\Delta). \tag{3.13}$$

On a first glance there are similarities with the continuum measure (1.55). Using $g(\xi) \sim A^2$ we see that the power of the "local areas" in both measures

equals $-3/2$. However, reparametrization invariance fixes the continuum measure completely, while in the discretized version (3.8) might be multiplied by the diffeomorphism invariant area factor $A^{1-2\beta/3}$, leading to the measure

$$d\mu(l_1,\dots,l_L) = \text{const} \times \frac{\sqrt{P(A_1^{\frac{2\beta}{3}},\dots,A_T^{\frac{2\beta}{3}})}}{\prod_{k=1}^{T} A_k^{\beta}} \prod_{j=1}^{L} l_j dl_j \; \delta(\Delta). \tag{3.14}$$

This generalization stresses the fact that the power of the areas is not fixed in quantum Regge calculus by any obvious principle since the areas are diffeomorphism invariant quantities with no direct local interpretation in the continuum.

The measure (3.14) is highly nonlocal and thus not suited for numerical simulations. Below we will show that it does not reproduce continuum physics.

3.2.2 The DeWitt-like measure in other dimensions than two

We can perform the derivation of the DeWitt-like measure in one and in higher dimensions. A one-dimensional piecewise linear manifold consists of L links of lengths l_i $(x_i = l_i^2), i = 1,\dots,L$, which are glued together at their ends. The canonical metric is thus given by $g^i_{\mu\nu} = (x_i)$. Therefore the DeWitt metric gives:

$$\langle \delta g, \delta g \rangle_T = (1 + \frac{C}{2}) \sum_{i=1}^{L} x_i^{-\frac{3}{2}} (\delta x_i)^2. \tag{3.15}$$

Thus the DeWitt-like measure in one dimension is

$$d\mu(l_1,\dots,l_L) = \text{const} \times \prod_{i=1}^{L} \frac{dl_i}{l_i^{\frac{1}{2}}}. \tag{3.16}$$

In this case the measure is local. The factor $(1 + \frac{Cd}{2})$ factorizes as in the continuum. Note that in $d = 1$ we cannot use the canonical value $C = -2$ for C since the DeWitt metric would be singular.

In higher dimensions the situation is more complicated. For a d-simplex the measure is derived from the scalar product

$$\langle \delta g, \delta g \rangle_T = \delta x M_d \delta x, \tag{3.17}$$

where δx is the $\frac{d(d+1)}{2}$-dimensional vector associated with link length deformations of the d-simplex. As in two dimensions, the complete norm for the

triangulation can be written as a straightforward superposition of the matrices M_d. However, in dimensions larger than two, the entries of this matrix M_d^{total} depend on the x_i's. In general, the measure is given by

$$d\mu(l_1, \ldots, l_L) = \text{const} \times \sqrt{\det M_D^{total}} \prod_{i=1}^{L} l_i dl_i \; \delta(\Delta), \tag{3.18}$$

where $\delta(\Delta)$ now stands for the generalized triangle inequalities. As an example we have computed the measure for the simplest closed three-dimensional manifold which consists of two tetrahedra glued together along the six links. The result is

$$d\mu(l_1, \ldots, l_6) = \text{const} \times \frac{1}{V} \prod_{i=1}^{6} l_i dl_i \; \delta(\Delta), \tag{3.19}$$

where V is the 3-volume of the manifold.

Note that also in four dimensions, where the DeWitt metric is simply $\prod_{\xi \in M} \prod_{\mu \leq \nu} dg_{\mu\nu}$, the determinant of M_d^{total} does not evaluate to a constant, not even for the simplest 4-geometries.

3.2.3 The DeWitt-like measures for special geometries

We have not been able to obtain a general closed expression for the polynomial P. However, in a number of special cases it is possible to find explicit expressions for the measure (3.13) which can immediately be generalized for the measure (3.14) [2].

The easiest case of a closed piecewise linear manifold are two triangles of area A glued together along the three links. The measure is

$$d\mu(l_1, l_2, l_3) = \text{const} \times \frac{1}{A^{\frac{3}{2}}} \prod_{j=1}^{3} l_j dl_j \; \delta(\Delta). \tag{3.20}$$

A tetrahedron has six links and four faces. The determinant of the 6×6 matrix M can be computed. The measure is given by

$$d\mu(l_1, \ldots, l_6) = \text{const} \times \frac{\sum_{i=1}^{4} A_i}{\prod_{i=1}^{4} A_i} \prod_{j=1}^{6} l_j dl_j \; \delta(\Delta). \tag{3.21}$$

Several other more complicated examples are possible. As a last example we display the measure for a hedgehog geometry which consists of K hat-like building blocks such as the one shown in figure 3.1. This block consists of



Figure 3.1: Building blocks for the hedgehog geometry which consists of K of these blocks glued together along the links l_{a_k} and l_{b_k}.

two triangles with areas A_{2k-1} and A_{2k} respectively, glued together along the links l_{2k-1} and l_{2k}. The hedgehog geometry is built by gluing these elements together along the links l_{b_k} and $l_{a_{k+1}}$ successively. After this gluing the vertices x and y have the order $2K$, while the other vertices are of order two. Although this geometry might seem a bit artificial it plays an important role in the analysis of the phase transitions of two dimensional quantum gravity coupled to matter fields [69]. The matrix M for this geometry can be written in a block diagonal form. Its determinant can be computed, resulting in the measure

$$\begin{aligned} d\mu(l_1,\ldots,l_{3K}) &= \text{const} \times \frac{\prod_{j=1}^{3K} l_j dl_j}{\prod_{k=1}^{2K} A_k^{3/2}} \prod_{k=1}^{K} (A_{2k-1}+A_{2k})^{1/2} \\ &\quad \times \left| \prod_{k=1}^{K} A_{2k} + (-1)^{K-1} \prod_{k=1}^{K} A_{2k-1} \right| . \end{aligned} \tag{3.22}$$

For an even number of elements this measure vanishes whenever the product over the even areas equals the product over the odd areas. For odd K, the measure is always positive. Clearly, this points to a severe sickness of the measure (3.13).

3.2.4 Commonly used measures

As has been mentioned above, the DeWitt-like measures (3.14) are not suited for numerical simulations. Also, we have demonstrated, that the measure in quantum Regge calculus is not fixed as in the continuum case. Therefore it is natural to search for other, local measures for which a sensible continuum

limit can be defined. The principle of universality should ensure that all reasonable choices of the measure result in the same continuum theory.

For numerical simulations in two-dimensional quantum Regge calculus, a number of local measures have been suggested [87, 95, 96]. In all cases these are of the form

$$d\mu(l_1, \ldots, l_L) = \frac{1}{\prod_{j=1}^{T} A_j^{\beta}} \prod_{i=1}^{L} l_i^{-\alpha} dl_i \; \delta(\Delta), \tag{3.23}$$

for some parameter α or β, or derived from this[2]. These measures are arrived at by intuitively translating the continuum measure $\prod_{\xi \in M} \prod_{\mu \leq \nu} dg_{\mu\nu}$ as $\prod_{i=1}^{L} l_i dl_i$ and by replacing the product $\prod_{\xi \in M} g(\xi)^{\sigma}$ by the product of the "local" areas $\prod_{k=1}^{T} A_k^{2\sigma}$. The most popular choices are the scale invariant measure $\prod_{i=1}^{L} \frac{dl_i}{l_i}$ and the uniform measure $\prod_{i=1}^{L} dx_i \sim \prod_{i=1}^{L} l_i dl_i$.

Motivation for the first of these measures is often suggested by an analogy to the scale invariant continuum-measure

$$\mathcal{D}g_{\mu\nu} = \prod_{x \in M} g(x)^{-\frac{d+1}{2}} \prod_{\mu \leq \nu} dg_{\mu\nu}(x), \tag{3.24}$$

which has been advocated for four-dimensional quantum gravity in [97, 98]. However, this measure is not diffeomorphism invariant, as has been stressed in section 1.2.1, although the opposite has been claimed and "proved" by wrong formal arguments [98, 99]. The DeWitt measure is the only diffeomorphism invariant measure for continuum quantum gravity. Therefore the measure (3.24) cannot replace the DeWitt measure in continuum quantum gravity and any translations to discretized versions should be avoided.

Also the DeWitt-like measure (3.13) cannot be used to motivate (3.23) in dimensions higher than one, since (3.13) is highly nonlocal while the measures (3.23) are local. Note that also in four dimensions, where the DeWitt measure is simply

$$\prod_{x \in M} \prod_{\mu \leq \nu} dg_{\mu\nu}(x), \tag{3.25}$$

the discrete DeWitt-like measure is very complicated and highly nonlocal and does not equal the uniform measure (3.23) with $\alpha = -1$ and $\beta = 0$, not even for the simplest 4-geometries. This clearly shows that translations like

$$\prod_{x \in M} g(x)^{\sigma} \prod_{\mu \leq \nu} dg_{\mu\nu}(x) \to \prod_{j=1}^{T} A_j^{2\sigma} \prod_{i=1}^{L} l_i dl_i \; \delta(\Delta) \tag{3.26}$$

[2]Sometimes the product over all triangle areas is replaced by the product over all areas assigned to the links. In our arguments this is not important and corresponds to choosing another value for the parameters α and β.

or similar are far too simple minded, since the Jacobian of this variable transformation is not taken into account. The correct treatment of the diffeomorphisms and the Jacobian leads to the results in section 3.4.2.

However, it has been argued that, appealing to universality, the precise form of the measure need not to be known. In the continuum limit, averages of observables should result in their continuum values independent of details of the discretization like the exponents α and β in (3.23) or the underlying connectivity of the piecewise linear space. But numerical simulations revealed that averages of observables in the context of quantum Regge calculus

- do not reproduce results of continuum quantum gravity. In two dimensions not even the KPZ exponents (1.14) could be obtained.
- show a significant dependence on the measure.
- show a significant dependence on the underlying triangulation.

These are serious flaws in the Regge approach to quantum gravity.

3.2.5 Numerical simulations in two-dimensional quantum Regge calculus

Quantum Regge calculus has repeatedly been used for numerical simulations in two, three and four dimensions. However, its status as a theory of quantum gravity remained unclear even in two dimensions. To clarify this situation, two groups have recently performed large scale numerical investigations in two dimensions where it is possible to compare with continuum results.

Bock and Vink [91] have performed a Monte Carlo simulation of two-dimensional Regge calculus applied to pure gravity. They added the term

$$\beta \int_M d^2\xi \sqrt{g}\, \mathcal{R}^2 \tag{3.27}$$

to the action. Such curvature square terms have been introduced in the context of four-dimensional quantum gravity to overcome the unboundedness of the Einstein-Hilbert action from below, which is induced by conformal fluctuations [100]. In two dimensions this term is the only nontrivial part of the action. Its discretized version can be written as

$$\beta \sum_{i=1}^{V} \frac{\delta_i^2}{A_{(i)}}, \tag{3.28}$$

h	0 (sphere)	1 (torus)	2 (bitorus)
γ in Liouville theory	-0.5	2	4.5
γ in QRC from [91]	$\gtrapprox 5.5$	≈ 2.0	$\gtrapprox 5.5$

Table 3.1: Values for the susceptibility exponent γ for pure two-dimensional gravity at different topologies as computed in Liouville theory compared to the values from the Monte Carlo simulation in [91]. Bock and Vink do not give error bars for their values. However, the deviation from the Liouville values is significant.

where δ_i is the deficit angle at the vertex i and $A_{(i)}$ is the area assigned to the vertex i by a barycentric division of the triangle areas around i. The addition of this term to the action allows a scaling analysis which gives the susceptibility exponent γ. This exponent can for example be defined as an entropy exponent for the partition function

$$Z(V) \propto V^{3-\gamma} e^{\Lambda_c V}, \tag{3.29}$$

where the dominant exponential is nonuniversal and the universal part is given by the subleading power correction. Continuum calculations show, that for pure gravity γ is given by

$$\gamma = 2 - \frac{5}{2}(1-h), \tag{3.30}$$

compare (1.14). It has been shown in [71, 101], that the addition of a small curvature square term does not change the value of γ in dynamical triangulation.

To be conservative, Bock and Vink used the scale invariant measure (3.23) with $\alpha = 1$ and $\beta = 0$ for their simulations. This enabled them to compare their results with previous calculations [96], where agreement between quantum Regge calculus and the formula (3.30) has been reported. They computed the susceptibility exponent for the topology of a sphere ($h = 0$), a torus ($h = 1$) and a bitorus, that means a sphere with two handles ($h = 2$). The results are compared with the continuum values in table 3.1. For spherical and bitoroidal topology they found that γ in quantum Regge calculus differs significantly from Liouville theory. That means that quantum Regge calculus does not reproduce the fundamental KPZ-result (3.30). Note that the torus topology is not well suited to test the KPZ formula (1.14). For $h = 1$ γ is independent of the conformal charge of the matter coupled to

gravity. Furthermore γ takes the same value 2 in the classical case without metrical fluctuations ($c = -\infty$) and in the quantum case.

The results of Bock and Vink have first been confirmed [102] and later been questioned [103, 104] by Holm and Janke. Using a very careful finite size scaling ansatz for pure quantum gravity on a fixed random triangulation of the sphere, they arrive at the value $\gamma = -10(2)$ for the susceptibility exponent [93], in the same setup. This is in clear disagreement with the prediction from Liouville theory and with the computations in [91, 96].

Using an alternative measure similar to (3.23) with $\alpha = 0$ and $\beta = \frac{1}{2}$ it was found in [102] that a scaling can no longer be observed and one encounters numerical difficulties. The observables did not reach equilibrium values. A similar change in the scaling behaviour was observed in [91]. The value of γ is changed when the scale invariant measure is replaced by $\prod_{i=1}^{L} \frac{dl_i}{l_i} l_i^{\zeta}$, for $\zeta \neq 0$. Furthermore it has been demonstrated in [93], that the results of quantum Regge calculus depend on the chosen fixed triangulation.

3.3 The appearance of spikes

In this section we will show, that two-dimensional quantum Regge calculus does not reproduce results of continuum quantum gravity with any of the proposed measures mentioned above [2].

Let us recall that the partition function $G(V, R)$ for closed two-dimensional universes with fixed volume V and with two marked points separated a geodesic distance R has the asymptotic behaviour (1.33)

$$G(V, R) \sim V^{-1/4} e^{-\left(\frac{R}{V^{1/4}}\right)^{\frac{4}{3}}}, \quad \text{for } \frac{R}{V^{1/4}} \to \infty. \tag{3.31}$$

Thus the expectation value of any power of the radius R can be calculated as

$$\langle R^n \rangle = \int_0^{\infty} dr\ r^n G(V, r) \sim V^{n/4}. \tag{3.32}$$

We can show that (3.32) is not fulfilled in quantum Regge calculus.

The physical reason is, that contrary to continuum quantum gravity in two dimensions, the volume, or equivalently the cosmological constant, does *not* set an intrinsic scale for the theory. It follows that the concept of length has no natural definition in this formalism and that every generic manifold degenerates into spikes.

3.3.1 General proof of appearance

Theorem 1 *For any value of α or β in (3.14) and (3.23) there exists an n such that for any link l in a given arbitrary triangulation*

$$\langle l^n \rangle_V = \infty \tag{3.33}$$

for any fixed value V of the spacetime volume. Thus the average radius $\langle R \rangle$, or some suitable power $\langle R^n \rangle$, does not exist [2].

Proof: In the situation of figure 3.2, which displays a small part of the piece-



Figure 3.2: Parametrization of the link lengths and the triangle areas in a part of the piecewise linear manifold M. The vertex v is of order N. A spike can be formed by making the links $l_1, \ldots, l_N$ arbitrary large. When the area of the surface is fixed, the link lengths $l_{N+1}, \ldots, l_{2N}$ have to be small of order $1/l_1$.

wise linear manifold M, the vertex v has the arbitrary order N. The links connected to v are labelled as $l_1, \ldots, l_N$, while the links opposite to v are labelled as $l_{N+1}, \ldots, l_{2N}$. We want to analyze the Regge integral in that part of the phase space where the vertex v is pulled to infinity and forms a spike while the volume of M is held bounded. In this situation the link lengths $l_1, \ldots, l_N$ are large, while the link lengths $l_{N+1}, \ldots, l_{2N}$ have to be of the order l_1^{-1}.

First we analyze the measure (3.23) with $\beta = 0$. Let Λ be a large number. Then l_1 can be integrated freely between Λ and infinity, while the

integration over $l_2, \dots, l_N$ is restricted by triangle inequalities. The integral over $l_2, \dots, l_N$ gives a factor $l_1^{(1-N)\alpha}(l_{N+1} \dots l_{2N})^{\frac{N-1}{N}}$ after symmetrizing over $l_{N+1}, \dots, l_{2N}$. An additional factor $l_{N+1} \dots l_{2N}$ comes from the triangle inequalities for the adjacent triangles which contain the links $l_{N+1}, \dots, l_{2N}$. Taking these factors together results in

$$\int_\Lambda^\infty dl_1 \; l_1^{-N\alpha} \int_0^{\frac{\Lambda}{l_1}} dl_{N+1} \dots dl_{2N} \; (l_{N+1} \dots l_{2N})^{\frac{2N-1}{N}-\alpha}, \tag{3.34}$$

which exists if

$$\alpha < 3 - \frac{1}{N}. \tag{3.35}$$

If that is fulfilled, the integral over $l_{N+1} \dots l_{2N}$ can be performed, which gives

$$\int_\Lambda^\infty dl_1 \; l_1^{-N\alpha} \left(\frac{\Lambda}{l_1}\right)^{N\left(\frac{3N-1}{N}-\alpha\right)} \sim \int_\Lambda^\infty dl_1 \; l_1^{1-3N}. \tag{3.36}$$

This means that $\langle l_1^n \rangle = \infty$ for all $n \geq 3N-2$, which proves the theorem 1 for the measure (3.23) with $\beta = 0$..

To discuss the general form of (3.23) we have to parametrize the areas $A_1, \dots, A_{2N}$. In our situation these are given as the products of a small and a large link length if ν is pulled to infinity. Thus the product $\prod_{j=1}^T A_j^{-\beta}$ gives an extra factor $(l_1 \dots l_N)^{-\beta}(l_{N+1} \dots l_{2N})^{-2\beta}$. After integrating over $l_2, \dots, l_N$ and after taking the triangle inequalities into account we get:

$$\int_\Lambda^\infty dl_1 \; l_1^{-N(\alpha+\beta)} \int_0^{\frac{\Lambda}{l_1}} dl_{N+1} \dots dl_{2N} \; (l_{N+1} \dots l_{2N})^{\frac{2N-1}{N}-\alpha-2\beta}, \tag{3.37}$$

which exists if $\alpha + 2\beta < 3 - \frac{1}{N}$. Then the integrals can be computed to give

$$\int_\Lambda^\infty dl_1 \; l_1^{1+N(\beta-3)}. \tag{3.38}$$

Thus for all $n \geq N(3-\beta) - 2$ the expectation value of l_1^n is infinite.

For the measure (3.14) the same analysis can be repeated if one notes, that from the nonlocal factor $\sqrt{P(A_1^{2\beta/3}, \dots, A_T^{2\beta/3})}$ the product $\prod_{i=1}^L l_i^{\beta/3}$ can be split of. It turns out that this doesn't change the result and we get again (3.38). This completes the proof. We denote $\langle l^n \rangle = \infty$ as the appearance of spikes.

3.3.2 Spikes in special geometries

Clearly, if a discretized theory shall make any sense, the results should not depend on details like the chosen triangulation. In fact however, the proof of theorem 1 shows, that $\langle l^n \rangle$ depends crucially on the order of the vertices, to which the link l is attached. To illustrate this dependence of quantum Regge calculus on the underlying triangulation further, we analyze some special geometries in the same way as in the proof to theorem 1. For some geometries even the partition function is ill defined for some measures, while for other measures or geometries $\langle l^n \rangle$ might be finite for some values of n.

Spikes in a hexagonal geometry



Figure 3.3: Parametrization of a hexagonal geometry; $k \neq m$. All centers of the cells, which are marked by thicker lines form spikes.

We analyze a regular triangulation with torus topology in which all vertices have the order 6, consisting of K hexagonal cells and parametrized as in figure 3.3. We want to study the measure (3.23) in that part of the configuration space, where all centers of cells form spikes while the total area of the surface is held bounded to prevent exponential damping from the action. That means that the 6K link lengths $l_i, i = 1, \ldots 6K$ are very large while the 3K link lengths $l_{a_k}, l_{b_k}, l_{c_k}, k = 0, \ldots, K-1$ are very small, of order l_1^{-1}. In cell number k one link, say l_{6k+1} can be integrated freely from a large number Λ to infinity, while the integration over $l_{6k+2}, \ldots, l_{6k+6}$ is then constrained by the triangle inequalities. Integrating these out and symmetrizing over the

small links $l_{a_k}, l_{b_k}, l_{c_k}, k = 0, \ldots, K-1$ gives

$$\int_\Lambda^\infty \prod_{k=1}^{K} l_{6k+1}^{-6(\alpha+\beta)} \int_0^{\frac{\Lambda}{l_{6k+1}}} \prod_{k=1}^{K} (l_{a_k} l_{b_k} l_{c_k})^{\frac{5}{3}-\alpha-2\beta} dl_{a_k} dl_{b_k} dl_{c_k}. \tag{3.39}$$

Thus the integral over $l_{a_k}, l_{b_k}, l_{c_k}$ only exists if $\alpha + 2\beta < \frac{8}{3}$. Furthermore $\langle l_1^n \rangle = \infty$ if $n \geq 7 + 3\alpha$.

For the measure (3.14) we extract a factor $\prod_{i=1}^{L} l_i^{\frac{\beta}{3}}$ from the nonlocal factor. Performing the analysis gives the bound $\beta < \frac{11}{5}$ on β. $\langle l_1^n \rangle$ is infinite if $n \geq 4 - \beta$. These results are summarized in table 3.2.

Spikes in a 12-3 geometry

Similarly we can analyze the integration measures for quantum Regge calculus for a triangulation of the torus where two thirds of the vertices have order 3 and one third of the vertices have order 12, see figure 3.4. We consider that



Figure 3.4: Illustration of the 12-3 geometry. All vertices of order 3 form spikes such that the total area of the surface is held bounded from above, to prevent exponential damping from the action.

part of the configuration space where all vertices of order 3 form spikes. The bounds on the exponents α and β in (3.23) and (3.14) are depicted in table 3.2. The resulting bounds are sharper because the order of the vertices which form spikes is lower which leaves less geometrical restriction. For $\alpha \leq -1$ the the average of a link length cannot be defined with the measure (3.23) (independent of β). The same holds true for the measure (3.14) if $\beta > 0$.

Spikes in degenerate geometries

One can get sharper bounds for geometries with vertices of order two, like the hedgehog geometry introduced in section 3.2.3. As explained in that section, these geometries do occur in phases of simplicial quantum gravity in the framework of dynamical triangulation and it would be unnatural to forbid their occurrence as basically local parts of very large triangulations.

As a first example we analyze the situation for the geometry shown in



Figure 3.5: A degenerate geometry which has two vertices of high order while all other vertices are of order 4. Spikes can be formed in several ways. In the text we analyze the behaviour in that part of the phase space where every second vertex of order 4 forms a spike.

figure 3.5 which has two vertices of high order while all other vertices are of order 4. This is slightly more regular than the hedgehog geometry associated with figure 3.1. In table 3.2 we present the results from our analysis of the situation, where every second vertex of order 4 forms a spike. While with measure (3.23) $\langle l^2 \rangle$ is undefined for $\alpha \leq -1$, the DeWitt-like measure (3.13) does not even admit the computation of the average $\langle l \rangle$.

For the hedgehog geometry these measures themselves are illdefined, see table 3.2. Measure (3.23) is illdefined for $\alpha = -1$ (the uniform measure), while for instance for the scale invariant measure (3.23) with $\alpha = 1$ and $\beta = 0$ expectation values of l^2 and higher are infinite.

Geometry	Measure	Bounds	Values of n s.t. $\langle l^n \rangle = \infty$
hexagonal,	(3.23)	$\alpha + 2\beta < \frac{8}{3}$, $-\frac{7}{3} < \alpha$	$n \geq 7 + 3\alpha$
see figure 3.3	(3.14)	$\beta < \frac{11}{5}$	$n \geq 4 - \beta$
12-3, see	(3.23)	$\alpha + 2\beta < \frac{7}{3}$, $-\frac{5}{3} < \alpha$	$n \geq \frac{5}{2} + \frac{3}{2}\alpha$
figure 3.4	(3.14)	$\beta < 2$	$n \geq 1 - \frac{\beta}{2}$
degenerate,	(3.23)	$\alpha + 2\beta < \frac{5}{2}$, $-2 < \alpha$	$n \geq 4 + 2\alpha$
see figure 3.5	(3.14)	$\beta < \frac{21}{10}$	$n \geq 2 - \frac{2\beta}{3}$
hedgehog, see	(3.23)	$\alpha + 2\beta < 2$, $-1 < \alpha$	$n \geq 1 + \alpha$
section 3.2.3	(3.14)	$\beta < 0$	$n \geq -\frac{\beta}{3}$

Table 3.2: Bounds for the exponents α and β for the measures (3.23) and (3.14) and values of n such that $\langle l^n \rangle = \infty$ for various geometries.

3.4 Concluding remarks

3.4.1 The appearance of spikes in higher dimensions

Although the derivation in the last section has been given only for the two-dimensional case, we expect, that one has to deal with the same problems in higher-dimensional quantum Regge calculus. The measures (3.23) have obvious generalizations in higher dimensions. An extra problem in higher dimensions is the unboundedness of the action which can be overcome by adding a cutoff involving for instance a curvature squared term. However, if the cutoff is taken to zero, we expect the spikes to reemerge. Actually, this has been observed in numerical simulations [105, 106]. In [106] four-dimensional quantum Regge calculus is explored on a fixed but not regular triangulation of the 4-torus. The triangulation is constructed from a regular triangulation with 31 vertices following [107] by inserting three additional vertices with low order. While in the conventional regular triangulation of the 4-torus all vertices are of order 30, the additional vertices are of order 5. The Einstein-Hilbert action plus a cosmological constant term is discretized in the usual way. A cutoff is introduced by the requirement that the local

fatness

$$\phi_s = 576 \frac{V_s^2}{\max_{l_i \in s}(l_i^8)} \tag{3.40}$$

for every 4-simplex s is held bounded away from zero, $\phi_s \geq f = \text{const} > 0$. Here we have introduced the 4-volume V_s of the simplex s. The fatness is maximal for equilateral simplices and vanishes for collapsing ones.

For the integration over the link lengths l_i the measure

$$d\mu(l_1, \ldots, l_L) = \prod_{i=1}^{L} l_i^{2\sigma-1} dl_i \delta(\Delta) \tag{3.41}$$

has been chosen for σ between 0 and 1. In figure 1 of [106] the expectation value of l_i^2 is depicted as a function of the cutoff f at the values $f = 2^m 10^{-6}$, $m = 9, \ldots, 0$ for $\sigma = 1$, $\sigma = \frac{1}{2}$ and $\sigma = \frac{1}{10}$. For $\sigma = 1$ and $\sigma = \frac{1}{2}$ this expectation value tends to infinity if the cutoff is removed while for $\sigma = \frac{1}{10}$ the situation cannot be decided from the numerical data in [106]. From the figure we deduce that the behaviour of $\langle l_i^2 \rangle$ is a power of f:

$$\langle l_i^2 \rangle \sim f^{-0.26(2)}, \text{ for } \sigma = 1, \tag{3.42}$$

$$\langle l_i^2 \rangle \sim f^{-0.16(1)}, \text{ for } \sigma = \frac{1}{2}. \tag{3.43}$$

This clearly demonstrates the appearance of spikes in four dimensions and shows that one has to expect the dependence of the physical results on the choice of the measure and on the choice of the fixed triangulation in four dimensions as well as in two dimensions.

3.4.2 The role of diffeomorphisms in two dimensions

While we have shown that quantum Regge calculus in its present form is not a suitable candidate for a theory of quantum gravity it is an interesting question, whether one can develop a theory of quantum gravity in the spirit of quantum Regge calculus.

A promising alternative approach to the functional integration (1.1) was studied in [108, 94, 109]. The integration over all metrics modulo diffeomorphisms is replaced by the continuum functional integral over piecewise linear metrics with a finite number of singularities modulo diffeomorphisms. The singularities are the V vertices of the piecewise linear space at which the curvature is located.

This approach is remarkable since at each stage in the calculations diffeomorphisms are treated exactly. Thus it includes all metrics which are related

to a piecewise linear metric by diffeomorphisms. It turns out, that this is sufficient to reproduce a discretized form of the Liouville action. Therefore it is believed, that for $V \to \infty$ this discretized version approaches the continuum expressions in some sense. No ad hoc assumption about the form of the measure has to be made. Note that the connectivity of the piecewise linear space is not fixed. In particular, all metrics of dynamical triangulation are contained in this approach.

We will give a short review of these ideas for the two-dimensional case. As in the continuum calculations, the starting point is the DeWitt metric. Since the reduction of the degrees of freedom involves only geometries but not diffeomorphisms, the gauge fixing which leads to (1.72) can be performed in the same way as in the continuum. Now the integration $\mathcal{D}\phi$ over the conformal factor has to be restricted to those conformal factors which describe Regge geometries. For spherical topology, to which we confine the discussion here, there are no Teichmüller parameters and one can choose a unique background metric $\hat{g}$. Menotti et al. adopt the usual way of a stereographic projection of the piecewise linear surface on the plane with $\hat{g}_{\mu\nu} = \delta_{\mu\nu}$ [110]. Then the conformal factor for piecewise linear geometries can be parametrized as

$$e^{\phi} = e^{2\lambda_0} \prod_{i=1}^{V} |z - z_i|^{2(\alpha_i - 1)}. \tag{3.44}$$

Here $2\pi\alpha_i$ is the angular aperture at the vertex i while the z_i are the coordinates of the singularities in the complex plane. The sum of the deficit angles $1 - \alpha_i$ has to equal the Euler characteristic,

$$\sum_{i=1}^{V} (1 - \alpha_i) = 2, \tag{3.45}$$

that means that α_V can be expressed by the other angles. λ_0 is an overall scale factor. Note that this parametrization simply stems from a generalization of the common Schwarz-Christoffel mappings which map the complex upper half plane to an arbitrary polygon [111]. One advantage of this parametrization over a parametrization via link lengths is that there are no triangle inequalities.

Due to the presence of six conformal Killing vectors, compare section 1.2.2, the gauge fixing is not complete and the conformal factor ϕ is invariant under an $SL(2, \mathbb{C})$ transformation given by

$$z' = \frac{az + b}{cz + d}, \;\; ad - bc = 1, \;\; a, b, c, d \in \mathbb{C}. \tag{3.46}$$

A short calculation gives

$$\phi'(z', \lambda_0, z_i, \alpha_i) = \phi(z', \lambda_0', z_i', \alpha_i), \tag{3.47}$$

with

$$\lambda_0' = \lambda_0 - \sum_{i=1}^{V} (1 - \alpha_i) \log |cz_i + d|. \tag{3.48}$$

Geometrical invariants as the angles α_i and the area

$$A = e^{2\lambda_0} \int d^2z \prod_{i=1}^{V} |z - z_i|^{2(\alpha_i - 1)} \tag{3.49}$$

are invariant under the transformation (3.46). A counting shows, that as in the usual parametrization in Regge calculus there are $3V - 6$ dynamical degrees of freedom.

To compute the Fadeev-Popov determinant for these conformal factors one employs the same techniques as in the continuum. Namely one computes the change of

$$\log \frac{\det'(P^+P)}{\det\langle \omega_a, \omega_b \rangle_V} \tag{3.50}$$

under a variation of the parameters λ_0, z_i and α_i and integrates the result back. One gets [94]:

$$\log \sqrt{\frac{\det'(P^+P)}{\det\langle \omega_a, \omega_b \rangle_V}} = \tag{3.51}$$
$$\frac{26}{12} \left\{ \sum_{i,j \neq i} \frac{(1 - \alpha_i)(1 - \alpha_j)}{\alpha_i} \log |z_i - z_j| + \lambda_0 \sum_i (\alpha_i - \frac{1}{\alpha_i}) - \sum_i F(\alpha_i) \right\}.$$

This is the discretized form of the Liouville action. In the limit $V \to \infty$ it approaches the continuum Liouville action. The function F depends only on the angles α_i. For $V \to \infty$, $\sum_{i=1}^{V} F(\alpha_i)$ goes to a topological invariant. The discretized Liouville action is invariant under the $SL(2, \mathbb{C})$ transformation (3.46). Indeed, it has been shown in [112] that this invariance induced by the conformal Killing vectors is sufficient to derive the form (3.51) of the Liouville action up to the sum over the $F(\alpha_i)$. Note that the discretized Liouville action is strongly unlocal. Overcounting of equivalent locally flat geometries is avoided by expression (3.51). Indeed, an angle $\alpha_i = 1$ simply does not contribute to (3.51).

The integration measure for the conformal factor is defined by the scalar product $ds^2 = \int d^2z\ e^{\phi}\ \delta\phi\delta\phi$. The change to the variables $(q_1, \ldots, q_{3V}) = ((z_1)_x, \ldots, (z_V)_y, \lambda_0, \alpha_1, \ldots, \alpha_{V-1})$ involves a Jacobian:

$$\mathcal{D}_{e^{\phi}\delta_{\mu\nu}}\phi = \prod_{i=1}^{V} d^2z_i \prod_{j=1}^{V-1} d\alpha_j d\lambda_0\ \sqrt{\det J}, \tag{3.52}$$

which is given by the determinant of

$$J_{ij} = \int d^2z\ e^{\phi} \frac{\partial\phi}{\partial q_i}\frac{\partial\phi}{\partial q_j}, \tag{3.53}$$

with the conformal factor ϕ defined by (3.44). The diagonal entries J_{ii} for $i = 1, \ldots, 2V$ of this matrix require some regularization. The explicit form of the measure is unfortunately unknown even for the simplest geometries which makes the approach presently inaccessible for numerical tests. While it has been possible to extract the dependence of the Jacobian on the scale factor λ_0, almost nothing is known about the other variables. It is an interesting open problem to show that the conformal anomaly arises from this measure.

3.4.3 Conclusion

We have shown that in two-dimensional quantum Regge calculus the expectation value of a suitable power of a single link length diverges. No natural scale is set by the measures (3.23) or (3.14). Even if the total area is kept bounded from above, the probability of having arbitrary large link lengths is not exponentially suppressed. In two-dimensional quantum gravity however, the probability of having two points separated a distance R is exponentially suppressed as $e^{-R\sqrt{\Lambda}}$ for a given cosmological constant Λ and suppressed as $e^{-R^{4/3}/V^{1/3}}$ if the area V of spacetime is fixed. That means, that quantum Regge calculus in two dimensions does not provide a formulation of quantum gravity. In addition, the mere existence of the partition function depends crucially on the chosen measure as well as on the chosen fixed triangulation. Clearly, this is a severe problem: if quantum Regge calculus was a realistic theory, these microscopic details should be unimportant and not affect physical expectation values.

Furthermore, we have shown that in four dimensions spikes have been observed in numerical studies of quantum Regge calculus. Numerical investigations demonstrate that also in four dimensions the results do depend crucially on the underlying triangulation and on the measure. Thus our conclusion in four dimensions is the same as in two dimensions: Quantum

Regge calculus does not provide a theory of quantum gravity. A sensible continuum limit cannot be defined. Not even the concept of length can be defined unambiguously.

Conclusion

In this work we have analyzed the fractal structure of two-dimensional quantum gravity. A central result is that the spectral dimension equals two for all types of conformal matter with central charge smaller than one coupled to gravity. Other aspects of the fractal nature of the spacetime have been introduced and discussed such as the extrinsic and intrinsic Hausdorff dimension, the branching ratio into minimal bottleneck baby universes and the scaling of the two-point function. We have compared our results with the latest numerical data. Numerical simulations are performed in the framework of dynamical triangulation. We have discussed this method and illustrated its power by a computation of an exact expression of the two-point function in two-dimensional quantum gravity. Furthermore we have discussed quantum Regge calculus which has been suggested as an alternative discretization of quantum gravity. We have shown that this theory does not provide a suitable discretization of quantum gravity since it disagrees with the continuum results. This point constitutes another central result of this work.

There are several open questions which have to be addressed in future work. It would be very interesting to develop a method which would allow the solution of Liouville theory. More modestly, the computation of the geodesic distance or similar geometrical objects would be interesting. The derivation of the intrinsic Hausdorff dimension is not very well understood. An alternative derivation of this result would provide further insight into the theory. Related to this is the question, why the fractal dimension seems to equal four for unitary matter coupled to quantum gravity in two dimensions. Of course the final triumph would be an analogous analysis of the fractal properties of the quantum spacetime of four dimensional quantum gravity in the continuum approach and in the discretized approach.

While this work settles the question about the viability of quantum Regge calculus in its present formulation it would be interesting to develop different schemes to discretize quantum gravity. We have discussed one possibility in chapter 3. It would be interesting to study further properties of this approach.

Bibliography

[1] J. Ambjørn, D. Boulatov, J. L. Nielsen, J. Rolf, and Y. Watabiki, "The spectral dimension of $2-$d quantum gravity," *JHEP 02(1998)010* hep-th/9801099.

[2] J. Ambjørn, J. L. Nielsen, J. Rolf, and G. Savvidy, "Spikes in quantum Regge calculus," *Class. Quant. Grav.* 14 (1997) 3225, gr-qc/9704079.

[3] S. W. Hawking, "Space-time foam," *Nucl. Phys.* B144 (1978) 349.

[4] S. W. Hawking, "The path integral approach to quantum gravity," in *General Relativity. An Einstein Centenary Survey*, S. W. Hawking and W. Israel, eds., p. 746. Cambridge University Press, 1979.

[5] C. J. Isham, "Structural issues in quantum gravity," in *General Relativity and Gravitation: GR14*, p. 167. World Scientific, Singapore, 1997. gr-qc/9510063.

[6] A. M. Polyakov, "Quantum geometry of bosonic strings," *Phys. Lett.* B103 (1981) 207.

[7] V. G. Knizhnik, A. M. Polyakov, and A. B. Zamolodchikov, "Fractal structure of $2-$d quantum gravity," *Mod. Phys. Lett.* A3 (1988) 819.

[8] F. David, "Conformal field theories coupled to $2-$d gravity in the conformal gauge," *Mod. Phys. Lett.* A3 (1988) 1651.

[9] J. Distler and H. Kawai, "Conformal field theory and $2-$d quantum gravity," *Nucl. Phys.* B321 (1989) 509.

[10] J. Ambjørn, J. L. Nielsen, and J. Rolf, "Two-dimensional quantum geometry," in *Proceedings of the 31st International Symposium Ahrenshoop on the Theory of Elementary Particles, Buckow, 1997.*

[11] P. Ginsparg and G. Moore, "Lectures on $2-$d gravity and $2-$d string theory," in *Recent Directions in Particle Theory: From Superstrings and Black Holes to the Standard Model (TASI - 92): proceedings*, J. Harvey and J. Polchinski, eds. World Scientific, 1993. hep-th/9304011.

[12] A. A. Belavin, A. M. Polyakov, and A. B. Zamolodchikov, "Infinite conformal symmetry in two-dimensional quantum field theory," *Nucl. Phys.* B241 (1984) 333.

[13] J. L. Cardy, "Conformal invariance," in *Phase Transitions and Critical Phenomena*, C. Domb and J. L. Lebowitz, eds., vol. 11, p. 55. Academic Press, 1987.

[14] J. L. Cardy, "Conformal invariance and statistical mechanics," in *Les Houches Summer School 1988*, p. 169. North-Holland, 1989.

[15] A. Einstein, "Hamiltonsches Prinzip und allgemeine Relativitätstheorie," *Preuss. Akad. Wiss. Berlin, Sitzber.* (1916) 1111.

[16] D. Hilbert, "Die Grundlagen der Physik," *Königl. Gesell. d. Wiss. Göttingen, Nachr. Math. Phys. Kl.* (1915) 395.

[17] M. G. Harris, "The branching of graphs in $2-\mathrm{d}$ quantum gravity," *Mod. Phys. Lett.* A11 (1996) 2351, hep-th/9607136.

[18] F. David, "A scenario for the $\mathrm{c} > 1$ barrier in noncritical bosonic strings," *Nucl. Phys.* B487 (1997) 633, hep-th/9610037.

[19] J. B. Hartle and S. W. Hawking, "Wave function of the universe," *Phys. Rev.* D28 (1983) 2960.

[20] D. Ruelle, *Statistical Mechanics*. W. A. Benjamin Inc., Reading, 1977.

[21] J. Ambjørn, B. Durhuus, and T. Jonsson, "Summing over all genera for $\mathrm{d} > 1$: A toy model," *Phys. Lett.* B244 (1990) 403.

[22] H. Kawai, N. Kawamoto, T. Mogami, and Y. Watabiki, "Transfer matrix formalism for two-dimensional quantum gravity and fractal structures of space-time," *Phys. Lett.* B306 (1993) 19, hep-th/9302133.

[23] Y. Watabiki, "Construction of noncritical string field theory by transfer matrix formalism in dynamical triangulation," *Nucl. Phys.* B441 (1995) 119, hep-th/9401096.

[24] E. Mottola, "Functional integration over geometries," *J. Math. Phys.* 36 (1995) 2470, hep-th/9502109.

[25] O. Alvarez, "Theory of strings with boundaries: Fluctuations, topology, and quantum geometry," *Nucl. Phys.* B216 (1983) 125.

[26] G. Moore and P. Nelson, "Measure for moduli," *Nucl. Phys.* B266 (1986) 58.

[27] B. S. DeWitt, "Quantum theory of gravity. I. The canonical theory," *Phys. Rev* 160 (1967) 1113.

[28] K. Fujikawa, "Path integral measure for gravitational interactions," *Nucl. Phys.* B226 (1983) 437.

[29] K. Fujikawa and O. Yasuda, "Path integral for gravity and supergravity," *Nucl. Phys.* B245 (1984) 436.

[30] L. S. Brown, "Stress-tensor trace anomaly in a gravitational metric: Scalar fields," *Phys. Rev.* D15 (1977) 1469.

[31] S. W. Hawking, "Zeta function regularization of path integrals in curved space-time," *Commun. Math. Phys.* 55 (1977) 133.

[32] J. Polchinski, "Evaluation of the one loop string path integral," *Commun. Math. Phys.* 104 (1986) 37.

[33] Y. Watabiki, "Analytic study of fractal structure of quantized surface in two-dimensional quantum gravity," *Prog. Theor. Phys. Suppl.* 114 (1993) 1.

[34] J. Distler, Z. Hlousek, and H. Kawai, "Hausdorff dimension of continuous Polyakov's random surfaces," *Int. J. Mod. Phys.* A5 (1990) 1093.

[35] H. Kawai, "Quantum gravity and random surfaces," *Nucl. Phys. B (Proc. Suppl.)* 26 (1992) 93.

[36] J. Jurkiewicz and A. Krzywicki, "On the size of a Polyakov surface," *Phys. Lett.* B148 (1984) 148.

[37] B. DeWitt, "Quantum gravity: the new synthesis," in *General Relativity. An Einstein Centenary Survey*, S. W. Hawking and W. Israel, eds., p. 680. Cambridge University Press, 1979.

[38] J. Ambjørn, K. Anagnostopoulos, T. Ichihara, L. Jensen, and Y. Watabiki, "Quantum geometry and diffusion." NBI-HE-97-54, TIT/HEP-352, 1998.

[39] J. Ambjørn, J. Jurkiewicz, and Y. Watabiki, "On the fractal structure of two-dimensional quantum gravity," *Nucl. Phys.* B454 (1995) 313, `hep-lat/9507014`.

[40] S. Havlin and D. Ben-Avraham, "Diffusion in disordered media," *Advances in Physics* 36 (1987) 695.

[41] G. Thorleifsson and B. Petersson, "Beyond the $c = 1$ barrier in two-dimensional quantum gravity," `hep-lat/9709072`.

[42] S. Alexander and R. Orbach *J. Phys. Lett.* 43 (1982) L625.

[43] M. E. Cates, "Statics and dynamics of polymeric fractals," *Phys. Rev. Lett.* 53 (1984) 926.

[44] T. Jonsson and J. F. Wheater, "The spectral dimension of the branched polymer phase of two-dimensional quantum gravity," *Nucl. Phys.* B515 (1998) 549, hep-lat/9710024.

[45] J. D. Correia and J. F. Wheater, "The spectral dimension of nongeneric branched polymer ensembles," *Phys. Lett.* B422 (1998) 76, hep-th/9712058.

[46] Y. Watabiki, "Fractal structure of space-time in two-dimensional quantum gravity," in *Frontiers in Quantum Field Theory in honor of the 60th birthday of Keiji Kikkawa*, p. 158. World Scientific, 1996. hep-th/9605185.

[47] B. DeWitt, *Dynamical Theory of Groups and Fields.* Gordon and Breach Inc., 1965.

[48] N. Ishibashi and H. Kawai, "String field theory of $c \leq 1$ noncritical strings," *Phys. Lett.* B322 (1994) 67, hep-th/9312047.

[49] N. Ishibashi and H. Kawai, "A background independent formulation of noncritical string theory," *Phys. Lett.* B352 (1995) 75, hep-th/9503134.

[50] J. Ambjørn, K. Anagnostopoulos, T. Ichihara, L. Jensen, N. Kawamoto, Y. Watabiki, and K. Yotsuji, "Quantum geometry of topological gravity," *Phys. Lett.* B397 (1997) 177, hep-lat/9611032.

[51] J. Ambjørn, K. Anagnostopoulos, T. Ichihara, L. Jensen, N. Kawamoto, Y. Watabiki, and K. Yotsuji, "The quantum space-time of $c = -2$ gravity.," *Nucl.Phys.* B511 (1998) 673, hep-lat/9706009.

[52] S. Catterall, G. Thorleifsson, M. Bowick, and V. John, "Scaling and the fractal geometry of two-dimensional quantum gravity," *Phys. Lett.* B354 (1995) 58, hep-lat/9504009.

[53] J. Ambjørn and K. N. Anagnostopoulos, "Quantum geometry of $2 - \mathrm{d}$ gravity coupled to unitary matter," *Nucl. Phys.* B497 (1997) 445, hep-lat/9701006.

[54] K. Anagnostopoulos, P. Bialas, and G. Thorleifsson, "The Ising model on a quenched ensemble of $c = -5$ gravity graphs," cond-mat/9804137.

[55] J. Ambjørn, K. N. Anagnostopoulos, and G. Thorleifsson, "The quantum space-time of $c > 0$ 2d gravity," *Nucl. Phys. B (Proc. Suppl.)* 63 (1998) 742, hep-lat/9709025.

[56] J. Ambjørn, B. Durhuus, and J. Fröhlich, "Diseases of triangulated random surface models, and possible cures," *Nucl. Phys.* B257 (1985) 433.

[57] F. David, "Planar diagrams, two-dimensional lattice gravity and surface models," *Nucl. Phys.* B257 (1985) 45.

[58] V. A. Kazakov, A. A. Migdal, and I. K. Kostov, "Critical properties of randomly triangulated planar random surfaces," *Phys. Lett.* B157 (1985) 295.

[59] T. Regge, "General relativity without coordinates," *Il Nuovo Cimento* XIX (1961) 558.

[60] D. Weingarten, "Pathological lattice field theory for interacting strings," *Phys. Lett.* B90 (1980) 280.

[61] D. Weingarten, "Euclidean quantum gravity on a lattice," *Nucl. Phys.* B210 (1982) 229.

[62] J. Ambjørn, B. Durhuus, and T. Jonsson, *Quantum Geometry.* Cambridge University Press, 1997.

[63] F. David, "Simplicial quantum gravity and random lattices," in *Gravitation and Quantizations*, B. Julia and J. Zinn-Justin, eds., p. 77. North-Holland, 1995. hep-th/9303127.

[64] J. Ambjørn, "Quantization of geometry," in *Fluctuating geometries in statistical mechanics and field theory*, F. David, P. Ginsparg, and J. Zinn-Justin, eds., p. 77. North-Holland, 1996. hep-th/9411179.

[65] J. Ambjørn, J. Jurkiewicz, and Y. Watabiki, "Dynamical triangulations, a gateway to quantum gravity?," *J. Math. Phys.* 36 (1995) 6299, hep-th/9503108.

[66] P. D. Francesco, P. Ginsparg, and J. Zinn-Justin, "$2-$d gravity and random matrices," *Phys. Rep.* 254 (1995) 1, hep-th/9306153.

[67] W. T. Tutte, "A census of planar triangulations," *Canadian J. of Math.* 14 (1962) 21.

[68] S. Jain and S. D. Mathur, "World sheet geometry and baby universes in $2-$d quantum gravity," *Phys. Lett.* B286 (1992) 239, hep-th/9204017.

[69] J. Ambjørn, B. Durhuus, J. Fröhlich, and P. Orland, "The appearance of critical dimensions in regulated string theories," *Nucl. Phys.* B270 (1986) 457.

[70] D. V. Boulatov and V. A. Kazakov, "The Ising model on random planar lattice: The structure of phase transition and the exact critical exponents," *Phys. Lett.* B186 (1987) 379.

[71] F. David, "A model of random surfaces with non-trivial critical behavior," *Nucl. Phys.* B257 (1985) 543.

[72] F. David, "Randomly triangulated surfaces in two-dimensions," *Phys. Lett.* B159 (1985) 303.

[73] V. A. Kazakov, "Ising model on a dynamical planar random lattice: Exact solution," *Phys. Lett.* A119 (1986) 140.

[74] J. Ambjørn and J. Greensite, "Nonperturbative calculation of correlators in $2-$d quantum gravity," *Phys. Lett.* B254 (1991) 66.

[75] J. Ambjørn, C. V. Johnson, and T. R. Morris, "Stochastic quantization vs. KdV flows in $2-$d quantum gravity," *Nucl. Phys.* B374 (1992) 496, hep-th/9109018.

[76] G. Akemann, "Higher genus correlators for the hermitian matrix model with multiple cuts," *Nucl. Phys.* B482 (1996) 403, hep-th/9606004.

[77] F. David, "Loop equations and nonperturbative effects in two-dimensional quantum gravity," *Mod. Phys. Lett.* A5 (1990) 1019.

[78] J. Ambjørn, L. Chekhov, and Y. Makeenko, "Higher genus correlators and W infinity from the Hermitean one matrix model," *Phys. Lett.* B282 (1992) 341, hep-th/9203009.

[79] J. Ambjørn, L. Chekhov, C. F. Kristjansen, and Y. Makeenko, "Matrix model calculations beyond the spherical limit," *Nucl. Phys.* B404 (1993) 127, hep-th/9302014.

[80] T. R. Morris, "Checkered surfaces and complex matrices," *Nucl. Phys.* B356 (1991) 703.

[81] J. Ambjørn, J. Jurkiewicz, and Y. M. Makeenko, "Multiloop correlators for two-dimensional quantum gravity," *Phys. Lett.* B251 (1990) 517.

[82] J. Ambjørn, C. F. Kristjansen, and Y. M. Makeenko, "Higher genus correlators for the complex matrix model," *Mod. Phys. Lett.* A7 (1992) 3187, hep-th/9207020.

[83] V. A. Kazakov, "The appearance of matter fields from quantum fluctuations of $2-$d gravity," *Mod. Phys. Lett.* A4 (1989) 2125.

[84] J. Ambjørn and Y. M. Makeenko, "Properties of loop equations for the Hermitean matrix model and for two-dimensional quantum gravity," *Mod. Phys. Lett.* A5 (1990) 1753.

[85] J. Ambjørn and Y. Watabiki, "Scaling in quantum gravity," *Nucl. Phys.* B445 (1995) 129, hep-th/9501049.

[86] J. Ambjørn, M. Carfora, and A. Marzuoli, *The Geometry of Dynamical Triangulations*, vol. m 50 of *LNP*. Springer, 1997.

[87] H. W. Hamber, "Simplicial quantum gravity," in *Critical phenomena, random systems, gauge theory*, K. Osterwalder and R. Stora, eds., p. 375. North-Holland, 1986.

[88] R. M. Williams and P. A. Tuckey, "Regge calculus: A bibliography and brief review," *Class. Quant. Grav.* 9 (1992) 1409.

[89] R. M. Williams, "Recent progress in Regge calculus," *Nucl. Phys. B (Proc. Suppl.)* 57 (1997) 73, gr-qc/9702006.

[90] H. W. Hamber and R. M. Williams, "Two-dimensional simplicial quantum gravity," *Nucl. Phys.* B267 (1986) 482.

[91] W. Bock and J. C. Vink, "Failure of the Regge approach in two-dimensional quantum gravity," *Nucl. Phys.* B438 (1995) 320, hep-lat/9406018.

[92] C. Holm and W. Janke, "The Ising transition in $2-$d simplicial quantum gravity: Can Regge calculus be right?," *Nucl. Phys. B (Proc. Suppl.)* 42 (1995) 725, hep-lat/9501004.

[93] C. Holm and W. Janke, "Fixed versus random triangulations in 2d simplicial Regge calculus," *Phys. Lett.* B390 (1997) 59, hep-lat/9608151.

[94] P. Menotti and P. P. Peirano, "Functional integration on two-dimensional Regge geometries," *Nucl. Phys.* B473 (1996) 426, hep-th/9602002.

[95] M. Bander, "Functional measure for lattice gravity," *Phys. Rev. Lett.* 57 (1986) 1825.

[96] M. Gross and H. W. Hamber, "Critical properties of two-dimensional simplicial quantum gravity," *Nucl. Phys.* B364 (1991) 703.

[97] C. Misner, "Feynman quantization of general relativity," *Rev. Mod. Phys.* 29 (1957) 497.

[98] L. D. Faddeev and V. N. Popov, "Covariant quantization of the gravitational field," *Sov. Phys.-Usp.* 16 (1974) 777.

[99] H. W. Hamber and R. M. Williams, "On the measure in simplicial gravity," hep-th/9708019.

[100] G. W. Gibbons, S. W. Hawking, and M. J. Perry, "Path integrals and the indefininteness of the gravitational action," *Nucl. Phys.* B138 (1978) 141.

[101] D. V. Boulatov, V. A. Kazakov, I. K. Kostov, and A. A. Migdal, "Analytical and numerical study of the model of dynamically triangulated random surfaces," *Nucl. Phys.* B275 (1986) 641.

[102] C. Holm and W. Janke, "Measure dependence of $2-$d simplicial quantum gravity," *Nucl. Phys. B (Proc. Suppl.)* 42 (1995) 722, hep-lat/9501005.

[103] C. Holm and W. Janke, "New results on $\gamma_{\rm str}$ in $2-$d quantum gravity using Regge calculus," *Nucl. Phys. B (Proc. Suppl.)* 47 (1996) 621, hep-lat/9509051.

[104] C. Holm and W. Janke, "Measuring the string susceptibility in 2d simplicial quantum gravity using the Regge approach," *Nucl. Phys.* B477 (1996) 465, hep-lat/9511029.

[105] W. Beirl, E. Gerstenmayer, and H. Markum, "Influence of the measure on simplicial quantum gravity in four-dimensions," *Phys. Rev. Lett.* 69 (1992) 713, hep-lat/9204010.

[106] W. Beirl, H. Markum, and J. Riedler, "Regge gravity on general triangulations," *Phys. Lett.* B341 (1994) 12, hep-lat/9407020.

[107] Kühnel and Lassmann *Discrete Comp. Geom.* 3 (1988) 169.

[108] P. Menotti and P. P. Peirano, "Faddeev-Popov determinant in two-dimensional Regge gravity," *Phys. Lett.* B353 (1995) 444, hep-th/9503181.

[109] P. Menotti and P. P. Peirano, "Diffeomorphism invariant measure for finite dimensional geometries," *Nucl. Phys.* B488 (1997) 719, hep-th/9607071.

[110] D. Förster, "Nonperturbative $\mathrm{d}=2$ Euclidean gravity without a lattice," *Nucl. Phys.* B283 (1987) 669.

[111] V. I. Smirnov, *A course of higher mathematics*. Pergamon, Oxford, 1964.

[112] P. Menotti, "Group theoretical derivation of Liouville action for Regge surfaces," gr-qc/9707023.